

QUANTUM COHOMOLOGY OF $[\mathbb{C}^N/\mu_r]$

AREND BAYER AND CHARLES CADMAN

ABSTRACT. We give a construction of the moduli space of stable maps to the classifying stack $B\mu_r$ of a cyclic group by a sequence of r -th root constructions on $\overline{M}_{0,n}$. We prove a closed formula for the total Chern class of μ_r -eigenspaces of the Hodge bundle, and thus of the obstruction bundle of the genus zero Gromov-Witten theory of stacks of the form $[\mathbb{C}^N/\mu_r]$.

We deduce linear recursions for all genus-zero Gromov-Witten invariants.

1. INTRODUCTION

This paper combines two notions of stable maps—twisted ([AV02]) and weighted ([Has03])—to produce a formula for the genus 0 Gromov-Witten invariants of $[\mathbb{C}^N/\mu_r]$. More precisely, we derive a formula for the equivariant Euler class of the obstruction bundle on $\overline{M}_{0,n}([\mathbb{C}^N/\mu_r])$ as a pull-back of classes on $\overline{M}_{0,n}$. Our definition of weighted twisted stable maps is ad hoc, applying only to genus 0 maps with target $[\mathbb{C}^N/\mu_r]$. Nevertheless, there is a notion of obstruction bundle on each of these spaces, and we have a wall-crossing formula relating their equivariant Euler classes. When all except one of the weights is small, the equivariant Euler class is easy to compute, so we deduce in this way our formula on $\overline{M}_{0,n}([\mathbb{C}^N/\mu_r])$.

The starting point of this work is our theorem 2.5.2, which provides an explicit construction of $\overline{M}_{0,n}([\mathbb{C}^N/\mu_r])$ from $\overline{M}_{0,n}$ via root constructions. This motivates the generalization to weighted stable maps. They are defined by applying root constructions to the space of weighted stable curves.

1.1. Introduction to $[\mathbb{C}^N/\mu_r]$. The quantum cohomology of stack quotients of the form $[\mathbb{C}^N/G]$ has recently seen a lot of interest due to the crepant resolution conjecture (see [BG06]). However, they are also natural objects of study by themselves: whenever a smooth N -dimensional stack X has a local orbifold chart $[U/G]$ where G acts with an isolated fixed point, part of the quantum cohomology of X will be described by the quantum cohomology of $[\mathbb{C}^N/G]$. Moreover, if X has a torus action which restricts to the natural torus action on the chart $[\mathbb{C}^N/G]$, then the equivariant quantum cohomology of $[\mathbb{C}^N/G]$ is relevant for computing the quantum cohomology of X via localization.

The Chen-Ruan orbifold cohomology of $[\mathbb{C}^N/\mu_r]$ has a natural basis h_e for $e \in \mu_r$. Consider the moduli space $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ of twisted stable maps

Date: January 29, 2009.

The second author was supported by NSF grant No. 0502170.

to the origin $B\mu_r \cong [0/\mu_r] \subset [\mathbb{C}^N/\mu_r]$ in the sense of [AV02], where the branching behavior at the i -th section is prescribed by $e_i \in \mu_r$. The non-trivial equivariant Gromov-Witten invariants of $[\mathbb{C}^N/\mu_r]$ are given by integrals over these moduli spaces. The normal bundle to the origin is \mathbb{C}^N , understood as a vector bundle on $B\mu_r$ via the given μ_r -action. If we write the universal curve as $\pi: \overline{C} \rightarrow \overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$, and the universal map as $f: \overline{C} \rightarrow B\mu_r$, the obstruction bundle of the moduli space is $R^1\pi_*f^*\mathbb{C}^N$. The Gromov-Witten invariant for h_{e_1}, \dots, h_{e_n} is typically¹ given by the integral of the equivariant Euler class (with respect to the canonical action of the N -dimensional torus \mathcal{T} on \mathbb{C}^N) of the obstruction bundle

$$(1.1.1) \quad \langle h_{e_1} \otimes \dots \otimes h_{e_n} \rangle^{[\mathbb{C}^N/\mu_r]} = \int_{\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)} e_{\mathcal{T}}([R^1\pi_*f^*\mathbb{C}^N]).$$

We call these integrals *generalized Hurwitz-Hodge integrals*, as the obstruction bundle is a direct sum of μ_r -eigenspaces of the dual of the Hodge bundle, where the moduli space is to be understood as a compactification of the Hurwitz space of μ_r -covers of \mathbb{P}^1 by *admissible covers*.

1.2. Our methods and results. The starting point of our work is the following explicit description of this moduli space of stable maps to $B\mu_r$ via the r -th root construction of [Cad07]. Given a divisor D on a scheme X , the r -th root construction $X_{D,r}$ is a stack over X that is isomorphic to X outside of D , but whose points over D are stacky with μ_r as automorphism group. For every proper subset $T \subset [n-1] := \{1, \dots, n-1\}$ having at least 2 elements, let r_T be the order $\prod_{i \in T} e_i$, and let $D^T \subset \overline{M}_{0,n}$ be the divisor consisting of curves having a node which separates the marking labels $1, \dots, n$ into T and $[n] \setminus T$.

Theorem. $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ is a μ_r -gerbe over the stack constructed from $\overline{M}_{0,n}$ by successively doing the r_T -th root construction at the boundary divisor $D^T \subset \overline{M}_{0,n}$ for all proper subsets $T \subset [n-1]$ having at least 2 elements.

We prove this in theorem 2.5.2, and we also give an explicit description of the universal curve and of the r -torsion line bundle defining the morphism to $B\mu_r$; see definitions 2.3.1 and 2.5.1. The root constructions along the boundary divisor introduce the additional automorphisms of curves with stacky nodes, called *ghost automorphisms*.

Now assume that μ_r is acting linearly on \mathbb{C}^N with weights w_1, \dots, w_N . To determine a formula for the Chern class of the obstruction bundle, we use a reduction guided by the notion of weighted stable curves in [Has03] and weighted stable maps in [MM08, AG08, BM06]. Weighted stable curves with n marked points depend on weight data a_1, \dots, a_n , and yield many birational models of the moduli space $\overline{M}_{0,n}$. Particular choices of weight data lead to an explicit presentation of the moduli space $\overline{M}_{0,n}$ by a series of blow-ups starting with \mathbb{P}^{n-3} , such that each intermediate blow-up step has an interpretation as a moduli space.

¹The formula needs an additional factor in case there is a coordinate direction on which every e_i acts trivially.

Motivated by this work and guided by theorem 2.5.2, we make an ad-hoc definition of a “moduli space of weighted stable maps to $B\mu_r$ ” in section 3.3. When the weights are chosen such that all fibers of the universal curve are irreducible, the obstruction bundle can easily be computed from general facts about the r -th root construction; we do this in section 3.4 for the weight data that gives a moduli space isomorphic to \mathbb{P}^{n-3} .

By a careful analysis of the wall-crossing for changing weights in section 4, we can lift this to a closed formula for the equivariant top Chern class in equation (1.1.1) for the standard (non-weighted) stable maps. We will now state this formula in the case of $N = 1$ and the standard representation of μ_r with weight one:

For $1 \leq i \leq n$, let $\delta_i \in [0, 1)$ be the age of e_i , i.e. $e^{2\pi i \delta_i} = e_i$. For all subsets $T \subset [n]$, let $\delta_T = \sum_{i \in T} \delta_i$. Let $\langle x \rangle$ denote the fractional part of x if x is not an integer, and let $\langle x \rangle = 1$ if x is an integer. For $T \subset [n-1]$, $|T| \geq 2$, let ψ_T be any class in $H^*(\overline{M}_{0,n})$ such that the restriction $D^T \cdot \psi_T$ is the ψ -class of the node over D^T on the component corresponding to T ; e.g. we can set $\psi_T := -\psi_n + \sum_{[n-1] \supsetneq S \supsetneq T} D^S$.

Theorem. *The equivariant Euler class of the obstruction bundle for $[\mathbb{C}/\mu_r]$ is given as*

$$e_{\mathcal{T}}([R^1 \pi_* f^* \mathbb{C}]) = \prod_{p=\langle \delta_{[n-1]} \rangle}^{\delta_{[n-1]}-1} (t - p\psi_n) \cdot \prod_{\substack{T \subset [n-1] \\ 2 \leq |T|}} \prod_{p=\langle \delta_T \rangle}^{\delta_T-1} \left(1 + \frac{pD^T}{t + p\psi_T} \right)$$

The case of different weight follows by adjusting the ages in the above formula, and the case of \mathbb{C}^N by multiplying the individual classes in $H^*(\overline{M}_{0,n})$; the full formula is given in theorem 5.1.1.

In the appendix, it is shown that this class can be expressed as a continuous, piece-wise analytic function from a real $(n-1)$ -dimensional torus to $H_T^*(\overline{M}_{0,n}, \mathbb{R})$ that encodes equivariant top Chern classes for all stacks $[\mathbb{C}^N/\mu_r]$ where N, n and the weights w_1, \dots, w_N are fixed, and r and e_1, \dots, e_n are arbitrary. See the discussion after the proof of Lemma A.1.1.

By a generalized inclusion-exclusion principle, the Chern class formula leads to linear recursions for all Gromov-Witten invariants of $[\mathbb{C}^N/\mu_r]$ by a sum over partitions, where every partition corresponds to a moduli space of comb curves. They are particularly nice for local Calabi-Yau 3-folds $[\mathbb{C}^3/\mu_r]$. We deduce an explicit formula for the non-equivariant invariants of $[\mathbb{C}^3/\mu_3]$. These invariants are the integrals in equation (1.1.1) for which $N = r = 3$, n is a multiple of 3, and all $e_i = e^{2\pi i/3}$. The recursion we discovered for these numbers is

$$\begin{aligned} \langle h_{\omega}^{\otimes n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_3]} &= (-1)^{n+1} \left(\left(\frac{n-4}{3} \right)! \right)^3 \frac{1}{3} \\ &+ \sum_{p=1}^{\frac{n-3}{3}} \sum_{\underline{m}} \frac{(-1)^{|\underline{m}|+1}}{|\text{Aut } \underline{m}|} \prod_{j=1}^k \left(\left(m_j - \frac{2}{3} \right)! \right)^3 M(n-1, \underline{m}) \langle h_{\omega}^{\otimes n-3p} \rangle_{0,n}^{[\mathbb{C}^3/\mu_3]}, \end{aligned}$$

where the second sum is over all partitions $\underline{m} = (m_1, \dots, m_k)$ of p , $M(n-1, \underline{m})$ is the multinomial coefficient

$$M(n-1, \underline{m}) = \binom{n-1}{3m_1+1, \dots, 3m_k+1, n-1-\sum_j(3m_j+1)},$$

and $x! = \prod_{p=\langle x \rangle}^x p$. The base case is $\langle h_\omega^{\otimes 3} \rangle_{0,3}^{[\mathbb{C}^3/\mu_3]} = 1/3$.

1.3. Relation to other work. The construction of the moduli space and the universal curve via r -th roots has been described locally by Abramovich in [Abr05, §3.5]. The global description along with the explicit description of the universal map to $B\mu_r$ seems to be new.

Our approach is to describe maps to $B\mu_r$ by r -torsion line bundles, which is particularly convenient in combination with the r -th root constructions. Of course, one can instead work more geometrically with cyclic covers, which is the point of view adopted in [ACV03, CC07].

The Gromov-Witten theory of $[\mathbb{C}^N/G]$ has recently generated a lot of interest due to the “crepant resolution conjecture”; we refer to [BG06] for an introduction to the conjecture and overview of the existing literature. When X is an orbifold that admits a crepant resolution $\pi: Y \rightarrow X$, Ruan first conjectured that the quantum cohomology rings of Y and X are isomorphic [Rua02], and suggested that the q -variables for π -exceptional divisor classes on Y need to be specialized to -1 [Rua06] to recover the orbifold cohomology ring of X . In [BG06], the authors extended this conjecture: their claim can be formulated as a local linear isomorphism between the Frobenius manifolds of the quantum cohomology of X and Y (after analytic continuation). This isomorphism does not respect the natural origins of the two Frobenius manifolds, which corresponds to Ruan’s specialization of q -variables. When the action of G on \mathbb{C}^N leaves the volume form invariant, the stacks of the form $[\mathbb{C}^N/G]$ yield many non-trivial test cases for the conjecture. In this form it is only expected to hold for orbifolds satisfying the strong Lefschetz theorem; a more general formulation can be found in [CCIT06, section 5].

The results so far have been obtained by the use of one the following two techniques: either a combination of localization computations and use of the WDVV-equations, or by using Tseng’s computation of the Chern *character* of the obstruction bundle and Givental’s framework for Gromov-Witten theory.

In [BGP08], the authors explicitly determine the genus-zero Gromov-Witten potential of $[\mathbb{C}^2/\mu_3]$ and verify the crepant resolution conjecture. In [BG06], the case $[\mathbb{C}^2/\mu_2]$ is derived from the Hodge integral computations of [FP00].

More generally, the case of A_n -singularities $[\mathbb{C}^2/\mu_{n+1}]$ is shown in [CCIT07a, CCIT07b] based on the Chern character computation. Various other results have been announced in [BG06]. For $[\mathbb{C}^3/\mu_3]$, part of the potential is computed in [CCIT07a], up to the problem of inversion of the “mirror map”. While their technique is completely different to ours, our results are surprisingly close, as explained in section 3.5 and 6.4; our recursion can be interpreted as a combinatorial inversion of the mirror map.

Different recursions for invariants of $[\mathbb{C}^3/\mu_3]$ have been established by the second author and R. Cavalieri in [CC07], using localization on the space of twisted stable maps to μ_3 -gerbes over \mathbb{P}^1 .

While our results are quite general, we make no attempt at verifying the crepant resolution conjecture.

1.4. Acknowledgements. This project owes special thanks to R. Cavalieri; his explanations of [BGP08] got us started on the direction of this work. The first author would like to thank A. Bertram, Y. Iwao, Y.-P. Lee and G. Todorov for discussions about Givental's formalism and [CCIT07a] that helped understanding the relation to our results.

1.5. Notations and conventions. We write $[n]$ for the set $[n] = \{1, 2, \dots, n\}$. We write $\langle x \rangle = x - [x] + 1 \in (0, 1]$ for the fractional part of x , set to 1 if x is integral. At various places we will write $\prod_{p=\langle x \rangle}^x f(p)$ for the product $\prod_{0 < p \leq x, \langle p \rangle = \langle x \rangle} f(p)$. If $x < 0$, the notation $\prod_{p=\langle x \rangle}^x f(p)$ means $\prod_{p=x+1}^{\langle x \rangle-1} \frac{1}{f(p)}$ (which is consistent with $\prod_{p=\langle x \rangle}^x f(p) = f(x) \cdot \prod_{p=\langle x \rangle}^{x-1} f(p)$ for all $x \in \mathbb{R}$).

For $x > 0$, we write $x!$ for the fractional factorial $x! = \prod_{p=\langle x \rangle}^x p$.

We identify the rational Chow groups of the moduli stacks of twisted stable maps $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ with that of its coarse moduli $\overline{M}_{0,n}$ via pull-back, and similarly for all other moduli stacks we construct. In the appendix, we introduce and explain a few non-standard notations for divisors on $\overline{M}_{0,n}$ that are particularly well-suited for our setting; most of it is only used in section 5.2, the exception being

$$\psi_T := -\psi_n + \sum_{[n-1] \supsetneq S \supsetneq T} D^S$$

for any $T \subset [n-1]$.

2. MODULI SPACE OF STABLE MAPS TO $B\mu_r$ VIA r -TH ROOTS

In this section, we show how to construct a component of the moduli space of genus zero stable maps to $B\mu_r$ from the moduli space $\overline{M}_{0,n}$ of stable curves of genus zero by a series of r -th root constructions.

2.1. User's guide to the r -th root construction. Given an effective Cartier divisor D of a Deligne-Mumford stack X , and a positive integer r which is invertible on X , the r -th root construction of [Cad07] produces a DM-stack $X_{D,r}$ with the following properties:

- (1) There is a canonical map $\pi: X_{D,r} \rightarrow X$ that is an isomorphism over $X \setminus D$.
- (2) Every point in $X_{D,r}$ lying over $D \subset X$ has stabilizer μ_r .
- (3) The preimage of D is an infinitesimal neighborhood of the μ_r -gerbe² \mathcal{D} over D parameterizing r -th roots of the fibers of $\mathcal{O}_X(D)|_D$: this is the

²A gerbe over D is a stack \mathcal{D} over D which étale locally admits a section and has the property that any two local sections are locally 2-isomorphic. A gerbe $\mathcal{D} \rightarrow D$ is a μ_r -gerbe if μ_r acts as the 2-automorphism group of every section in a compatible way.

stack whose objects are triples $(f: S \rightarrow D, L, \phi)$ where f is a morphism, L is a line bundle on S , and $\phi: L^r \rightarrow f^*\mathcal{O}_X(D)|_D$ is an isomorphism. (Only when $\mathcal{O}_X(D)|_D$ is the r -th power of a line bundle is \mathcal{D} isomorphic to $D \times B\mu_r$.)

- (4) On $X_{D,r}$, there is a line bundle $\mathcal{O}_{X_{D,r}}(\mathcal{D})$ with a section $s_{\mathcal{D}}$ and an isomorphism $\phi: \mathcal{O}(\mathcal{D})^r \rightarrow \pi^*(\mathcal{O}(D))$ such that $\phi(s_{\mathcal{D}}^r) = \pi^*(s_D)$. (Here $s_D \in \mathcal{O}(D)$ is the tautological section vanishing along D .)

The universality of the data in (4) is the defining property: giving a morphism $f: S \rightarrow X_{D,r}$ is equivalent to giving a quadruple (g, L, s, ϕ) where $g = \pi \circ f$ is a morphism to X , L is a line bundle on S , s a section, and $\phi: L^r \rightarrow g^*\mathcal{O}(D)$ is an isomorphism sending s^r to $g^*(s_D)$.

Locally, when $X = \text{Spec } A$ is affine and the divisor D is given by an equation $(x = 0)$, the r -th root construction is given by stack quotient $[(\text{Spec } A[u]/(u^r - x))/\mu_r]$ of the cyclic μ_r -cover branched at D , but of course globally such a cover may not exist.

To the best of our knowledge, the r -th root construction is originally due to A. Vistoli and spread as a rumor for quite some time. His notation is $\sqrt[r]{(X, D)}$.

We call $\mathcal{O}_{X_{D,r}}(\mathcal{D})$ the tautological line bundle of the r -th root construction at D . The zero stack $Z \subset X_{D,r}$ of $s_{\mathcal{D}}^r$ is the preimage of $D \subset X$. The zero stack of $s_{\mathcal{D}}$ is the gerbe \mathcal{D} . To simplify notation we write $\mathcal{O}(\frac{1}{r}D)$ to refer to $\mathcal{O}(\mathcal{D})$. More generally, if $d \in \frac{1}{r}\mathbb{Z}$, we write $\mathcal{O}(dD)$ for $\mathcal{O}(\mathcal{D})^{\otimes dr}$. This notation is particularly nice to describe the push-forward of line bundles along π :

$$(2.1.1) \quad \pi_*\mathcal{O}(dD) = \mathcal{O}(\lfloor d \rfloor D)$$

(This follows from [Cad07, Theorem 3.1.1].)

If X is an algebraic space, then the coarse moduli space of $X_{D,r}$ is X . When X is smooth and $D \subset X$ is smooth, then $X_{D,r}$ is smooth. The construction commutes with base change for a morphism $f: Y \rightarrow X$ such that $f^{-1}(D)$ is a Cartier divisor. (The construction can be generalized a little to make it compatible with arbitrary base change: see $X_{(L,s,r)}$ in [Cad07].)

If $\mathbb{D} = (D_1, \dots, D_n)$ is an n -tuple of Cartier divisors and $\vec{r} = (r_1, \dots, r_n)$, we can iterate the root constructions to obtain a stack denoted $X_{\mathbb{D}, \vec{r}}$. This stack can also be realized as the n -fold fiber product over X of the root stacks X_{D_i, r_i} . If X is smooth, each individual D_i is smooth and the D_i have normal crossing, then $X_{\mathbb{D}, \vec{r}}$ is smooth, too. If $d_i \in \frac{1}{r_i}\mathbb{Z}$, we extend the above notation by writing $\mathcal{O}(\sum_i d_i D_i)$ for the tensor product of the line bundles $\mathcal{O}(\mathcal{D}_i)^{d_i r_i}$, where $\mathcal{O}(\mathcal{D}_i)$ is the tautological bundle corresponding to the root construction along D_i .

It is instructive (and important for the construction of $\overline{M}_{0,n}(B\mu_r)$ later on) to compare the stacks $X_{\mathbb{D}, \vec{r}}$ and $X_{D,r}$ in the case where $n = 2$, $r = r_1 = r_2$, and $D = D_1 \cup D_2$. On $X_{\mathbb{D}, \vec{r}}$, the line bundle $\mathcal{O}(\frac{1}{r}(D_1 + D_2))$ with section $s_{\mathcal{D}_1} \cdot s_{\mathcal{D}_2}$ defines an r -th root of D , and thus there is a natural map

$$(2.1.2) \quad X_{\mathbb{D}, \vec{r}} \rightarrow X_{D,r}.$$

However, this is not an isomorphism if the divisors intersect, and one way to see this is by looking at stabilizer groups of points of the two stacks. If $x \in D_1 \cap D_2$, then the stabilizer group of the point in $X_{\mathbb{D}, \vec{r}}$ lying over x is $\mu_r \times \mu_r$. On the other hand, the stabilizer group of any point in the preimage of D in $X_{D, r}$ is μ_r . If X is smooth and D_1, D_2 are smooth with normal crossings, then these stacks are also distinguished by the fact that $X_{\mathbb{D}, \vec{r}}$ is smooth, while $X_{D, r}$ is singular over $D_1 \cap D_2$.

2.2. Stable maps to $B\mu_r$. Consider the moduli space $\overline{M}_{0,n}(B\mu_r)$ of balanced twisted stable maps of genus 0 to $B\mu_r$ in the sense of [AV02], where we work over \mathbb{C} . Such a map over a scheme S can be described by the following data:

- A stacky nodal curve \mathcal{C} over S , with n divisors $\Sigma_1, \dots, \Sigma_n$ in the smooth locus of \mathcal{C} , and
- a line bundle L on \mathcal{C} together with an isomorphism $\phi: L^r \rightarrow \mathcal{O}_{\mathcal{C}}$.

These have to satisfy various properties:

- The stacky curve \mathcal{C} is a scheme away from its nodes and the divisors Σ_i , and its nodes are *balanced*.
- Each Σ_i is a cyclotomic gerbe over S .
- If C is the coarse moduli space of \mathcal{C} , then the image of every divisor $\Sigma_i \subset \mathcal{C}$ in C is isomorphic to the image of a section $x_i: S \rightarrow C$, so that (C, x_1, \dots, x_n) becomes a stable curve of genus zero with n marked points.
- The map $\mathcal{C} \rightarrow B\mu_r$ induced by (L, ϕ) is representable.

The line bundle L is the pull-back of the line bundle on $B\mu_r$ given by the canonical one-dimensional representation of μ_r . Every point $x \in \Sigma_i$ has an automorphism group isomorphic to μ_p for some p dividing r . This identification is canonical if the representation of μ_p corresponding to the fiber of the normal bundle $\mathcal{O}_{\Sigma_i}(\Sigma_i)$ at x equals the standard representation. Let ω be the primitive r -th root of unity $\omega = e^{\frac{2\pi i}{r}}$. Then $\omega^{r/p}$ acts on the fiber L_{x_i} as multiplication by e_i for some $e_i \in \mu_r$. Equivalently, the map of stabilizer groups $\mu_p \rightarrow \mu_r$ (which is injective by representability of $\mathcal{C} \rightarrow B\mu_r$) sends the canonical generator of μ_p to e_i . These group elements e_1, \dots, e_n are constant on every connected component of $\overline{M}_{0,n}(B\mu_r)$.

From now on, we assume we are given $e_1, \dots, e_n \in \mu_r$ and restrict our attention to the connected component $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$. There is a natural map $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r) \rightarrow \overline{M}_{0,n}$ induced by the coarse moduli space of the universal curve. Our theorem will describe this map explicitly via a series of root constructions. One explanation for these root constructions is that twisted curves have ghost automorphisms for each twisted node (cf. [Abr05, section 3.5]).

2.3. Construction of the moduli space via root constructions. Let r and $e_i \in \mu_r, i = 1, \dots, n$ be given. For convenience we allow $e_i = 1$, i.e. untwisted points; then the universal curve is given by the forgetful morphism

$$\pi_{0,(e_1, \dots, e_n)}: \overline{M}_{0,n}(e_1, \dots, e_n, 1; B\mu_r) \rightarrow \overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r).$$

The component is empty unless $\prod_i e_i = 1$.

Consider the universal stable curve of genus zero $\pi_{0,n}: \overline{M}_{0,n+1} \rightarrow \overline{M}_{0,n}$. The boundary divisors of $\overline{M}_{0,n}$ are indexed by subsets $T \subset [n]$ such that $2 \leq |T| \leq n-2$ and $n \notin T$ (see Appendix A). For every such T , let r_T be the order of $\prod_{i \in T} e_i$.

Definition 2.3.1. Let $\overline{M}^{(1)}$ be the stack constructed from $\overline{M}_{0,n}$ by doing the r_T -th root construction at every boundary divisor D^T . We construct $\overline{C}^{(1)}$ from $\overline{M}_{0,n+1}$ in the same way after setting $e_{n+1} = 1$.

(In particular, we take the r_i -th root construction at every section $s_i = D^{i,n+1}$ where r_i is the order of e_i .)

Lemma 2.3.2. There is a canonical map $\pi^{(1)}: \overline{C}^{(1)} \rightarrow \overline{M}^{(1)}$.

Proof. Equivalently, we construct a map to the fiber product

$$\overline{C}^{(0)} = \overline{M}_{0,n+1} \times_{\overline{M}_{0,n}} \overline{M}^{(1)};$$

then $\overline{C}^{(0)}$ will be the relative coarse moduli space. Since $\pi_{0,n}^{-1}(D^T) = D^T \cup D^{T \cup \{n+1\}}$ and the r -th root construction is compatible with such a base change, this fiber product can be constructed from $\overline{M}_{0,n+1}$ by the r_T -th root constructions at all divisors $D^T \cup D^{T \cup \{n+1\}}$ for $T \subset [n]$. To construct $\overline{C}^{(1)}$, we instead took the r_T -th root construction at D^T and $D^{T \cup \{n+1\}}$ separately, hence the morphism $\overline{C}^{(1)} \rightarrow \overline{C}^{(0)}$ is given by forgetting the root construction along all sections, followed by a composition of morphisms as in (2.1.2) above. \square

Note that $\overline{C}^{(1)}$ has additional automorphisms along the nodes of the curves lying over D^T . When we restrict this family to the μ_{r_T} -gerbe in $\overline{M}^{(1)}$ lying over D^T , the fibers become stacky curves with a twisted node. The node is balanced because after base change to a scheme over the base, the remaining automorphism group is the kernel of the multiplication $\mu_{r_T} \times \mu_{r_T} \rightarrow \mu_{r_T}$, which acts with opposite weight on the two branches. The so-called “ghost automorphisms” are accounted for by the additional automorphism introduced in the moduli space. We have thus proved:

Proposition 2.3.3. The morphism $\pi^{(1)}: \overline{C}^{(1)} \rightarrow \overline{M}^{(1)}$ is a family of balanced twisted curves.

Note that we adapted [AV02, Definition 4.1.2] to a family over a Deligne-Mumford stack: all conditions have to be checked after étale base change to a scheme covering $\overline{M}^{(1)}$.

Each fiber of $\pi^{(1)}$ admits a morphism to $B\mu_r$ having the correct restrictions to Σ_i (given by e_1, \dots, e_n). However, these morphisms do not in general glue to a morphism $\overline{C}^{(1)} \rightarrow B\mu_r$. They will glue precisely when the μ_r -gerbe of Definition 2.5.1 is trivial.

2.4. The universal line bundle. For $T \subset [n]$, we will write always $T^C = [n] \setminus T$ for its complement.

Lemma 2.4.1. *Let C be a geometric fiber of $\pi^{(1)}$ and let*

$$\mathcal{L} = \mathcal{O}_{\overline{C}^{(1)}}\left(\frac{1}{r_T}D^{T,n+1}\right).$$

- (1) *If there is no node $x \in C$ such that one of the two connected components of $C \setminus \{x\}$ contains exactly the markings of T (and the other those of T^C), then $\mathcal{L}|_C$ is trivial.*
- (2) *Otherwise, let C_3, C_4 be the two connected components of C after normalization at x , such that C_3 contains all the markings of T , and C_4 those of T^C , and let $C_1 \subset C_3$ and $C_2 \subset C_4$ be the two irreducible components of C meeting at x . Then*

$$(2.4.1) \quad \mathcal{L}|_{C_1} \cong \mathcal{O}_{C_1}\left(-\frac{1}{r_T}x\right)$$

$$(2.4.2) \quad \mathcal{L}|_{C_2} \cong \mathcal{O}_{C_2}\left(\frac{1}{r_T}x\right)$$

$$(2.4.3) \quad \mathcal{L}|_{C'} \cong \mathcal{O}_{C'}$$

where C' is any irreducible component of C other than C_1, C_2 .

Proof. The first statement is obvious as C does not meet the divisor D^T in that case.

In the second case, $C_3 = C \cap D^{T,n+1}$, and so equation (2.4.2) is obvious, as is (2.4.3) for all $C' \subset C_4$. The claim then follows by symmetry and the fact that the restriction of $\mathcal{O}(\frac{1}{r_T}(D^T + D^{T,n+1}))$ to C is trivial, as it is the pull-back of the tautological line bundle $\mathcal{O}(\frac{1}{r_T}D^T)$ on $\overline{M}^{(1)}$. \square

Now choose $d_i \in \frac{1}{r} \cdot \mathbb{Z}$ such that $e^{2\pi i d_i} = e_i$ and $\sum_{i=1}^n d_i = 0$ (which is possible since $\prod_{i=1}^n e_i = 1$). For $T \subset [n]$, let $d_T = \sum_{i \in T} d_i$.

Lemma 2.4.2. *Define the line bundle L_1 on $\overline{C}^{(1)}$ as*

$$(2.4.4) \quad L_1 := \mathcal{O}\left(\sum_{i=1}^n d_i s_i + \sum_{\substack{T \subset [n] \\ 2 \leq |T| \leq n-2 \\ n \notin T}} d_T D^{T,n+1}\right)$$

Then L_1^r is the pull-back of a line bundle on $\overline{M}^{(1)}$:

$$L_1^r = (\pi^{(1)})^*(L_2)$$

We write $n \notin T$ to stress that this is for now just an arbitrary way to pick exactly one of T, T^C for all subsets T .

Proof. First note that L_1^r is pulled back from the coarse moduli space $\overline{M}_{0,n+1}$ since $d_T \in \frac{1}{r} \cdot \mathbb{Z}$ for all T . A line bundle on a family of nodal curves of genus zero over a smooth scheme is pulled back from the base if and only if its degree on any irreducible component of every fiber is zero (in which case it is the pull-back of its own push-forward to the base). Hence it is sufficient to check that the degree of L_1^r (or, equivalently, the degree of L_1) is zero on any irreducible component C_0 of any fiber C of π^1 (in which case L_1^r is even pulled back from $\overline{M}_{0,n}$).

Let x_1, \dots, x_m be the nodes of C contained in C_0 . Let $T_i \subset \{1, \dots, n\}$, $1 \leq i \leq m$ be the markings contained in the irreducible components which are connected to C_0 via the node x_i , and let T_0 be the markings contained in C_0 . For every j with $1 \leq j \leq m$ exactly one of $\mathcal{O}(d_{T_j} D^{T_j, n+1})$ and $\mathcal{O}(d_{T_j^C} D^{T_j^C, n+1})$ will appear in the right-hand side of equation (2.4.4) defining L_1 ; by the previous lemma and $d_T = -d_{T^C}$, both restrict to $\mathcal{O}(d_{T_j} x_j)$ on C_0 . By the same lemma, all other $\mathcal{O}(d_T D^{T, n+1})$ restrict trivially to C_0 . Hence

$$L_1|_{C_0} = \mathcal{O}\left(\sum_{i \in T_0} d_i s_i + \sum_{1 \leq j \leq m} d_{T_j} x_j\right).$$

This line bundle has degree $\sum_{j=0}^m \sum_{i \in T_j} d_i = 0$ since $[n]$ is the disjoint union of all T_j . \square

2.5. Base change to the gerbe.

Definition 2.5.1. Let $\overline{M}^{(2)}$ be the μ_r -gerbe over $\overline{M}^{(1)}$ of r -th roots of L_2 . Let $\pi^{(2)}: \overline{C}^{(2)} \rightarrow \overline{M}^{(2)}$ be the base change of $\pi^{(1)}: \overline{C}^{(1)} \rightarrow \overline{M}^{(1)}$ via $\overline{M}^{(2)} \rightarrow \overline{M}^{(1)}$, and let $L_2^{1/r}$ be the universal line bundle that is an r -th root of L_2 .

By abuse of notation, we write L_1 also for the pull-back of L_1 to $\overline{C}^{(2)}$. The line bundle $L = L_1 \otimes \pi^{(2)*} L_2^{-1/r}$ together with the obvious isomorphism $L^r \rightarrow \mathcal{O}_{\overline{C}^{(2)}}$ defines a morphism $\overline{C}^{(2)} \rightarrow B\mu_r$.

Theorem 2.5.2. *The following diagram*

$$\begin{array}{ccc} \overline{C}^{(2)} & \longrightarrow & B\mu_r \\ \pi^{(2)} \downarrow & & \\ \overline{M}^{(2)} & & \end{array}$$

is a family of twisted stable maps over $\overline{M}^{(2)}$ which defines an isomorphism

$$m: \overline{M}^{(2)} \rightarrow \overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r).$$

Proof. We already showed that $\pi^{(1)}$ (and thus $\pi^{(2)}$) is a family of balanced twisted curves.

The morphism $\overline{C}^{(2)} \rightarrow \overline{M}^{(2)} \times B\mu_r$ is representable: away from the sections and nodes, the map $\pi^{(2)}$ is already representable, and since all nodes and sections do not intersect each other, we can treat them separately. At a section s_i , the relative inertia group of $\pi^{(2)}$ is isomorphic to $B\mu_{r_i}$; since that group acts faithfully on L_1 and thus on L , the map on inertia groups is injective. A similar argument holds for all nodes.

We thus get a morphism m as claimed in the theorem. By lemma 2.5.3, m is an isomorphism if both stacks are smooth and the morphism is birational and a bijection of \mathbb{C} -valued points which induces isomorphisms of their stabilizer groups. Since $\overline{M}^{(2)}$ is étale over $\overline{M}^{(1)}$, which is a root construction on $\overline{M}_{0,n}$ at smooth

divisors with transversal intersection, it follows that $\overline{M}^{(2)}$ is smooth. The first order deformations of an n -marked, genus 0 twisted stable map to $B\mu_r$ are the same as those of the marked twisted curve, which has dimension equal to $n - 3$ (see [ACV03, section 3]). As this equals the dimension of $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$, it is also smooth.

For bijectivity, it suffices to show for each n -marked genus 0 curve C , there is a unique twisted stable map to $B\mu_r$ with coarse moduli space C and contact types e_1, \dots, e_n . For C irreducible, this uniqueness is shown in [CC08, 2.1.5]. Otherwise, one can show by induction that the contact types at the nodes are uniquely determined by those at marked points. So the morphism is unique over each component of C , and it suffices to show that it glues uniquely over the nodes. Since the morphism to $B\mu_r$ is equivalent to a line bundle L and a non-vanishing section of L^r , the gluing is clearly unique up to isomorphism.

It remains only to check the map on automorphism groups induced by m . If x is a closed point of $\overline{M}^{(2)}$, the automorphism group $G_{m(x)}$ of $m(x)$ in the moduli stack $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ can be identified with the group of μ_r -automorphisms of the μ_r -cover \tilde{C}_x of the fiber C_x of $\overline{C}^{(2)}$ over x . If S is the set of irreducible components of C_x , this identifies $G_{m(x)}$ with the subgroup of μ_r^S that acts compatibly over every node; since the preimage of a node $n_T \in C_x$ in \tilde{C}_x is isomorphic to μ_r/μ_{r_T} , we can identify $G_{m(x)}$ with the kernel of the map

$$\Sigma = \prod_T \Sigma_T: \mu_r^S \rightarrow \prod_{T \subset [n-1] | x \in D^T} \mu_r/\mu_{r_T}$$

where Σ_T is given by the quotient of the group elements corresponding to the two irreducible components meeting in n_T .

The μ_r -cover \tilde{C}_x is given by the r -th roots of unity inside the line bundle $L|_{C_x}$; hence to understand the map $G_x \rightarrow G_{m(x)}$ it is sufficient to look at how the automorphism group G_x of x acts on L . By the construction of $\overline{M}^{(2)}$, the automorphism group of x is

$$G_x = \mu_r \times \prod_{T \subset [n-1] | x \in D^T} \mu_{r_T}.$$

By the definition of L , the first factor acts via L_2 and thus diagonally, whereas μ_{r_T} acts diagonally on the irreducible components of $D^{T,n+1} \cap C_x$ (and trivially on all others), by its induced action on $\mathcal{O}(D^{T,n+1})$. Let C_0 be the component of C_x which contains the n -th marking.

Before showing that $m: G_x \rightarrow G_{m(x)}$ is an isomorphism, we introduce some notation. For $g \in G_x$, write g_0 for the projection of g onto μ_r , and write g_T for its projection onto μ_{r_T} . For each irreducible component C_i of C_x , let $C_{j_0}, C_{j_1}, C_{j_2}, \dots, C_{j_{k_i}}$ be the unique shortest path from C_i to C_0 . That is to say, $j_0 = i$, $j_{k_i} = 0$, C_{j_ℓ} meets $C_{j_{\ell+1}}$ in a node for each $0 \leq \ell \leq k_i - 1$, and there are no repetitions in j_0, \dots, j_{k_i} . For $0 \leq \ell \leq k_i - 1$, let T_ℓ be the subset of $[n-1]$ determined by the node joining

C_{j_ℓ} to $C_{j_{\ell+1}}$. Then $m(g)$ acts on the restriction of L to the component C_i by

$$(2.5.1) \quad g_0 \prod_{\ell=0}^{k_i-1} g_{T_\ell}.$$

Let $g \in G_x$, and suppose that $m(g)$ is trivial. Since $m(g)$ acts on C_0 by g_0 , it follows that g_0 is trivial. By inducting on the number of nodes separating a given node from C_0 , and using (2.5.1), it follows that g_T is trivial for each T . Therefore, g is trivial.

Now suppose $h \in G_{m(x)}$. Let g_0 be the element of μ_r by which h acts on C_0 . By inducting over the nodes as in the previous paragraph, and using the fact that the irreducible components of C_x form a tree, we can now define g_T for each T in such a way that $m(g) = h$. Therefore, $m : G_x \rightarrow G_{m(x)}$ is an isomorphism. \square

We remark that [ACV03, section 7] contains a careful treatment of automorphism groups for G -covers.

Lemma 2.5.3. *Let \mathcal{X} and \mathcal{Y} be normal, separated, integral, Deligne-Mumford stacks of finite type over an algebraically closed field k of characteristic zero. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a birational morphism which induces an equivalence of categories between objects over $\text{Spec } k$. Then f is an isomorphism.*

Proof. Let $V \rightarrow \mathcal{Y}$ be an étale surjective morphism from a scheme V and let $U = V \times_{\mathcal{Y}} \mathcal{X}$. Then $U \rightarrow V$ is separated and quasi-finite, hence quasi-affine by [LMB00, A.2]. Therefore, U is a scheme. Let $U' \subseteq U$ be a connected component and let $V' \subseteq V$ be its image. The hypotheses of the lemma imply that U' and V' are normal varieties and that $U' \rightarrow V'$ is a birational morphism which is bijective on k -points. By Zariski's birational correspondence theorem, it follows that $U' \rightarrow V'$ is an isomorphism. Applying the argument to each connected component shows that $U \rightarrow V$ is an isomorphism. It now follows from [LMB00, 3.8.1] that $\mathcal{X} \rightarrow \mathcal{Y}$ is an isomorphism. \square

2.6. Comments on the construction. The pull-back of a one-dimensional $B\mu_r$ -representation is a power of L . Hence, in order to understand the Chern class of the obstruction bundle $R^1\pi_*\mathbb{C}^N$ it is sufficient to understand the Chern classes of the higher direct image $R^1\pi_*L^w$ of powers of L , and their products.

It is worth pointing out that while the ghost automorphism groups are isomorphic to μ_{r_T} , this isomorphism is not natural; the ghost automorphism group is naturally isomorphic to the relative stabilizer group of the twisted node, and by choosing one of the two components $D^{T,n+1}$ or $D^{T^C,n+1}$ (and identifying the stabilizer group by its action on the corresponding tangent bundle) one gets an isomorphism to μ_{r_T} whose sign depends on this choice. In our construction, this choice shows up in the definition of L_1 , for which we had to choose one of T and T^C for all divisors D^T of $\overline{M}_{0,n}$. As L_2 depends on that choice only up to an r -th power, neither $\overline{M}^{(2)}$ nor the universal line bundle L depend on this choice. The map $m : \overline{M}^{(2)} \rightarrow \overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ does depend on it, however. Different

choices can be related by a composition with a ghost automorphism³ of the moduli stack.

3. WEIGHTED STABLE MAPS TO $B\mu_r$

3.1. Weighted stable maps. Let g be a genus, and $\mathcal{A} = (a_1, \dots, a_n)$ be *weight data*, which means $a_i \in \mathbb{Q} \cap [0, 1]$ satisfy $2g - 2 + \sum_i a_i > 0$.

In [Has03], Hassett introduced the notion of weighted stable curves: a weighted stable curve of type (g, \mathcal{A}) over S is a nodal curve $\pi: C \rightarrow S$ with n sections $s_i: S \rightarrow C$ such that

- (1) every section s_i with positive weight a_i is contained in the smooth locus of π ,
- (2) the rational divisor $K_{C/S} + \sum_i a_i s_i$ is π -relatively ample, and
- (3) for any $I \subset [n]$ such that the intersection $\bigcap_i s_i$ is non-empty, we have $\sum_i a_i \leq 1$.

We will summarize a few of his results, and refer to [Has03] for details.

If $a_i = 1$ for all i , then these are stable curves in the usual sense. The difference is that when points $s_i, i \in I$, collide, then only when $\sum_i a_i > 1$ does a new rational component bubble off. This is enough to make the new rational component stable according to condition (2).

All the moduli spaces $\overline{M}_{g, \mathcal{A}}$ with $|\mathcal{A}| = n$ are birational. More precisely, assume that the weight data $\mathcal{A} = (a_1, \dots, a_n)$ and $\mathcal{B} = (b_1, \dots, b_n)$ satisfy $a_i \geq b_i$ for all i , and $a_i > b_i$ for at least one i (we will write $\mathcal{A} > \mathcal{B}$ from now on). Then there is a birational reduction morphism $\rho_{\mathcal{B}, \mathcal{A}}: \overline{M}_{g, \mathcal{A}} \rightarrow \overline{M}_{g, \mathcal{B}}$. (It is induced by \mathcal{B} -stabilizing the family of curves over $\overline{M}_{g, \mathcal{A}}$.) There is a chamber decomposition of $[0, 1]^n$ by a finite number of walls such that the moduli space $\overline{M}_{g, \mathcal{A}}$ only depends on the chamber in which the weight data \mathcal{A} lies: the walls are associated to subsets $T \subset [n]$ and given as

$$(3.1.1) \quad w_T = \left\{ a_i \mid \sum_{i \in T} a_i = 1 \right\}.$$

Further, the contraction morphism for crossing a single wall is given as a smooth blow-up.

It is somewhat convenient to allow at least one weight to be zero, because $\overline{M}_{g, \mathcal{A} \cup \{0\}}$ is *by definition* the universal curve over $\overline{M}_{g, \mathcal{A}}$.

This notion has been extend to weighted stable maps in [MM08], [BM06] and [AG08]. In particular, in [BM06] and [AG08] it was shown that Gromov-Witten invariants can be computed for any choice of weights, yielding identical GW-invariants, and [AG08] gave wall-crossing formulae for the full Gromov-Witten potential including gravitational descendants.

3.2. $\overline{M}_{0, n}$ as a blow-up of \mathbb{P}^{n-3} . As an example that will be important later, consider for given n the weights $\mathcal{A}_k = (\frac{1}{k}, \dots, \frac{1}{k}, 1)$ (with $n - 1$ entries of $\frac{1}{k}$) for $k = 1, \dots, n - 2$. The moduli space $\overline{M}_{0, \mathcal{A}_{n-2}}$ is isomorphic to \mathbb{P}^{n-3} , and the universal curve is the blow-up $\text{Bl}_x \mathbb{P}^{n-2}$ of \mathbb{P}^{n-2} at a point x ; the universal map

³by which we mean an automorphism covering the identity on the coarse moduli space

is the projection of \mathbb{P}^{n-2} to $\mathbb{P}^{n-3} \cong \mathbb{P}T_x$ from x . If we pick x away from the coordinate hyperplanes, then the image of the special section with weight 1 is the exceptional divisor, while the remaining sections can be given as the coordinate hyperplanes. (The special section cannot intersect with any other, while the only condition on the remaining sections is that they may not *all* coincide.)

The moduli space $\overline{M}_{0,\mathcal{A}_{n-3}}$ is the blow-up of $\overline{M}_{0,\mathcal{A}_{n-2}}$ at the $n-1$ points that are the images of the intersections of $n-2$ of the $n-1$ coordinate hyperplanes. When we successively increase the first $n-1$ weights from $\frac{1}{n-2}$ to 1, one gets a description of $\overline{M}_{0,n}$ by successive blow-ups from \mathbb{P}^{n-3} . This is also explained in [Has03, section 6.2]; the description of $\overline{M}_{0,n}$ as a blow-up of \mathbb{P}^{n-3} is equivalent to the description by De Concini and Procesi in [DCP95].

3.3. The moduli spaces of weighted stable maps to $B\mu_r$ via r -th roots. We sidestep the question of defining a moduli problem of weighted stable maps to a stack in general. Instead we give a direct construction of the moduli stacks via r -th root constructions, guided by the construction in the non-weighted case in section 2.

Given r , weight data $\mathcal{A} > 0$, and $\mathcal{E} = (e_1, \dots, e_n) \in \mu_r^n$, we want to construct a stack which would resemble $\overline{M}_{0,\mathcal{A}}(e_1, \dots, e_n; B\mu_r)$ if it were to exist. Choose $d_i \in \frac{1}{r}\mathbb{Z}$ with $e^{2\pi i d_i} = e_i$ and $\sum_{i=1}^n d_i = 0$ as before. Boundary divisors on $\overline{M}_{0,\mathcal{A}}$ are given as D_σ for \mathcal{A} -stable 2-partitions $\sigma = (T, T^C)$ of $[n]$; \mathcal{A} -stable means that the condition $|T|, |T^C| \geq 2$ is replaced by $\sum_{i \in T} a_i > 1$ and $\sum_{i \in T^C} a_i > 1$. (This of course means that a corresponding rational curve with two components is \mathcal{A} -stable.)

Let $\overline{M}_{0,\mathcal{A}}^{(1)}$ be the stack obtained from $\overline{M}_{0,\mathcal{A}}$ by taking the r_T -th root at every divisor D^T such that (T, T^C) is \mathcal{A} -stable (where r_T is defined as before as the order of $\prod_{i \in T} e_i$). To obtain $\overline{C}_{0,\mathcal{A}}^{(1)}$ from $\overline{C}_{0,\mathcal{A}} = \overline{M}_{0,\mathcal{A} \cup \{0\}}$, we start with the same construction, but additionally construct the r_i -th root at every section s_i .⁴ The same proof as in lemma 2.3.2 shows:

Lemma 3.3.1. *There is a canonical map $\overline{C}_{0,\mathcal{A}}^{(1)} \rightarrow \overline{M}_{0,\mathcal{A}}^{(1)}$.*

This is a stacky curve with balanced nodes, but it can have points (in the relative smooth locus of the coarse moduli space) with automorphism group $\mu_r^{|I|}$, for curves where $s_i, i \in I$, are identical; so this is not a twisted stable curve in the sense of [AV02].

Proposition 3.3.2. *Let $L_{1,\mathcal{A}}$ be the line bundle on $\overline{C}^{(1)}$ defined by*

$$(3.3.1) \quad L_{1,\mathcal{A}} := \bigotimes_{i=1}^n \mathcal{O}(s_i)^{d_i} \otimes \bigotimes_T \mathcal{O}(D^{T,n+1})^{d_T}$$

where the second tensor product goes over all subsets $T \subset [n]$ with $n \notin T$ such that (T, T^C) is \mathcal{A} -stable. Then $L_{1,\mathcal{A}}^r = \pi^* L_{2,\mathcal{A}}$ for some line bundle $L_{2,\mathcal{A}}$ on $\overline{M}^{(1)}$.

⁴In the case of $\overline{M}_{0,n}$, the section s_i is equivalent to the boundary divisor given by $T = \{i, n+1\}$; however, this does not yield an \mathcal{A} -stable 2-partition, hence we need to list them separately.

Again, this has the same proof as before.

Let $\overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r}$ be the μ_r -gerbe over $\overline{M}_{0,\mathcal{A}}^{(1)}$ of r -th roots of $L_{2,\mathcal{A}}$, and let $\overline{C}_{0,\mathcal{A}}^{\mu_r}$ be the base change of $\overline{C}_{0,\mathcal{A}}^{(1)}$ to $\overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r}$. The line bundle $L_{\mathcal{A}} := L_{1,\mathcal{A}} \otimes \pi^* L_{2,\mathcal{A}}^{-1/r}$ on $\overline{C}_{\mathcal{A}}$ has trivial r -th power and thus defines a map $f: \overline{C}_{0,\mathcal{A}}^{\mu_r} \rightarrow B\mu_r$.

However, again $f: \overline{C}_{0,\mathcal{A}}^{\mu_r} \rightarrow B\mu_r$ is not a twisted stable map in the sense of [AV02]; most importantly, f is not representable (not representable even after a base change to a scheme $S \rightarrow \overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r}$).

3.4. \mathbb{P}^{n-3} -weight data. For any weight w of a one-dimensional μ_r -representation, we call

$$H_{\mathcal{A}}^w = R^1 \pi_* L_{\mathcal{A}}^w$$

the generalized dual Hodge bundle on $\overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r}$ for the weight data \mathcal{A} .

Our computation of the Chern class builds up from a direct computation for the weight data $\mathcal{A} = (\frac{1}{n-2}, \dots, \frac{1}{n-2}, 1)$, which yields $\overline{M}_{0,\mathcal{A}} \cong \mathbb{P}^{n-3}$ (see section 3.1). Given w , let $\delta_i^w \in [0, 1)$ be the age of the line bundle L^w at the i -th section; it is determined by $e^{2\pi i \delta_i^w} = e_i^w$. For any subset $T \subset [n]$, we let $\delta_T^w = \sum_{i \in T} \delta_i^w$.

Proposition 3.4.1. *The generalized dual Hodge bundle $H_{\mathcal{A}}^w$ has the following class in the K -group:⁵*

$$[H_{\mathcal{A}}^w] = \sum_{p=\langle \delta_{[n-1]}^w \rangle}^{\delta_{[n-1]}^w - 1} [\mathcal{O}(-pH)]$$

Proof. The moduli space is a μ_r -gerbe over \mathbb{P}^{n-3} , and the universal curve is constructed from $Bl_x \mathbb{P}^{n-2}$ by the r_i -th root construction at the section s_i for all i , and the base change to the μ_r -gerbe.

We choose d_i such that $d_i \in [0, 1)$ for $i = 1, \dots, n-1$ and $d_n = -\sum_{i=1}^{n-1} d_i$. Then $L_{2,\mathcal{A}}$ can be computed by $L_2 = s_n^*(L_{1,\mathcal{A}})^r = \mathcal{O}(-rd_n)$, and so by projection formula $H_{\mathcal{A}}^w = \mathcal{O}(wd_n) \otimes R^1 \pi_* L_{1,\mathcal{A}}^w$. To compute the higher direct image of $L_{1,\mathcal{A}}^w$, we break up π into the composition $\pi = \pi_2 \circ \pi_1$ of the map π_1 , forgetting the roots along the sections, with the map π_2 that is the base change of the natural projection $Bl_x \mathbb{P}^{n-2} \rightarrow \mathbb{P}^{n-3}$ to the μ_r -gerbe.

The push-forward along π_1 follows easily from equation (2.1.1). Using $-wd_n = \delta_{[n-1]}^w + \sum_{i=1}^{n-1} [wd_i]$, we get (where we write E for the exceptional divisor of $Bl_x \mathbb{P}^{n-2}$):

$$(\pi_1)_* L_{1,\mathcal{A}}^w = \mathcal{O} \left(\sum_{i=1}^n [wd_i] s_i \right) = \mathcal{O} \left((-wd_n - \delta_{[n-1]}^w) H + [wd_n] E \right)$$

⁵See 1.5 for other notation conventions used in this formula.

The relative canonical bundle of π_2 is $-H - E$, and the pull-back of the hyperplane class on \mathbb{P}^{n-3} is $H - E$. Hence

$$\begin{aligned} R^1(\pi_2)_*(\pi_1)_*L_{1,\mathcal{A}}^w &= \left((\pi_2)_* \mathcal{O} \left((wd_n + \delta_{[n-1]}^w - 1)H + (-\lfloor wd_n \rfloor - 1)E \right) \right)^\vee \\ &= \mathcal{O}(-\langle \delta_{[n-1]}^w \rangle - wd_n) \otimes \left((\pi_2)_* \mathcal{O} \left((\lceil \delta_{[n-1]}^w \rceil - 2)H \right) \right)^\vee, \end{aligned}$$

where we used $wd_n - \lfloor wd_n \rfloor = \{wd_n\} = 1 - \langle -wd_n \rangle = 1 - \langle \delta_{[n-1]}^w \rangle$ and $\delta_{[n-1]}^w - \langle \delta_{[n-1]}^w \rangle = \lceil \delta_{[n-1]}^w \rceil - 1$. Applying $(\pi_2)_*$ to the short exact sequences $\mathcal{O}((a-1)H) \rightarrow \mathcal{O}(aH) \rightarrow \mathcal{O}(aH)|_H$, and using $(\pi_2)_*\mathcal{O} = \mathcal{O}$, implies that in the K -group,

$$[R^1(\pi_2)_*(\pi_1)_*L_{1,\mathcal{A}}^w] = \sum_{p=0}^{\lceil \delta_{[n-1]}^w \rceil - 2} [\mathcal{O}(-\langle \delta_{[n-1]}^w \rangle - wd_n - p)].$$

Tensoring this with $\mathcal{O}(wd_n)$ and re-indexing yields the statement of the proposition. \square

Since the hyperplane class of \mathbb{P}^{n-3} agrees with the ψ -class of the n -th marking (which is special by having weight one), this implies:

Corollary 3.4.2. *In the situation of the previous proposition, the Chern class of $H_{\mathcal{A}}^2$ is given as*

$$c(H_{\mathcal{A}}^w) = \prod_{p=\langle \delta_{[n-1]}^w \rangle}^{\delta_{[n-1]}^w - 1} (1 - p\psi_n)$$

(Note that ψ_n denotes the pull-back of the corresponding class in $\overline{M}_{0,\mathcal{A}}$ by our convention for the rational Chow groups of the moduli stacks.)

3.5. Relation to the twisted I -function. In Givental's formalism for Gromov-Witten theory [Giv01, Giv04, CG07, CCIT07a], the so-called J -function plays an essential role. Let $X = [\mathbb{C}^N / \mu_r]$, where μ_r acts diagonally with weights w_1, \dots, w_N . The orbifold cohomology of X is $H = H^*(X) = \bigoplus_{e \in \mu_r} \mathbb{C} \cdot h_e$, and the J function is a map $H \rightarrow H[[z]][[z^{-1}]]$ defined by the following formula:

$$\begin{aligned} J^X(z, t) &= z + t + \sum_{n \geq 0} \sum_{e \in \mu_r} \frac{1}{n!} \langle t, \dots, t, \frac{h_e}{z - \psi} \rangle_{0, n+1}^X \cdot r h_{e^{-1}} \\ (3.5.1) \quad &= z + t + \sum_{n \geq 0} \sum_{e \in \mu_r} \sum_{k \geq 0} \frac{1}{n! z^{k+1}} \langle t, \dots, t, \psi_n^k h_e \rangle_{0, n+1}^X \cdot r h_{e^{-1}} \end{aligned}$$

For example, by the results of [JK02], the J -function of $B\mu_r$ is given as

$$J^{B\mu_r}(z, t) = z + t + \sum_{\underline{k}=(k_0, k_1, \dots, k_{r-1})} \prod_{j=0}^{r-1} \frac{t_j^{k_j}}{k_j! z^{|k_j|}} \cdot h_{\prod_j (\omega^j)^{k_j}}$$

where we wrote $t = \sum_{j=0}^{r-1} t_j h_{\omega^j}$. The idea of [CCIT07a] (and Givental's formalism in general), specialized to our setting, is to determine the J -function X from the J -function of $B\mu_r$; the former is called the *twisted J -function* in [CCIT07a].

As an approximation to the twisted J -function, Coates, Corti, Iritani and Tseng define a *twisted I -function* in [CCIT07a, section 4]; specialized to our case, and translated into our notation, it is given by⁶

$$I^{\text{tw}}(z, t) = z + t + \sum_{(e_1, \dots, e_n)} J^{(e_1, \dots, e_n)} \cdot M_{(e_1, \dots, e_n)}(z).$$

Here

$$J^{(e_1, \dots, e_n)} = z^{-n+2} \delta_{1, \prod_i e_i} \cdot h_{e_n}^{-1}$$

is the part of the J -function of $B\mu_r$ that comes from invariants computed on $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$, and $M_{(e_1, \dots, e_n)}(z)$ is defined by

$$M_{(e_1, \dots, e_n)}(z) = \prod_{a=1}^N \prod_{p=\langle \delta_{[n-1]}^{w_a} \rangle}^{\delta_{[n-1]}^{w_a}-1} (1 - pz).$$

To show that I^{tw} has the desired properties⁷ they use Tseng's Grothendieck-Riemann-Roch-computation of the Chern character of the obstruction bundle in [Tse05].

We can define a *weighted J -function* of X by

$$J^{X; \text{weighted}}(z, t) = z + t + \sum_{n \geq 0} \sum_{e \in \mu_r} \frac{1}{n!} \langle t, \dots, t, \frac{h_e}{z - \psi} \rangle_{0, n+1}^{X; \text{weighted}} \cdot r h_{e^{-1}},$$

where the invariant with superscript “weighted” denotes the invariant computed by the moduli of weighted stable maps $\overline{M}_{0, \mathcal{A}, \mathcal{E}}^{\mu_r}$ considered in the previous section, i.e. for $\mathcal{A} = (\frac{1}{n-2}, \dots, \frac{1}{n-2}, 1)$. Then the result of the previous section can be formulated as

$$J^{X; \text{weighted}}(z, t) = z + t + \sum_{(e_1, \dots, e_n)} J^{(e_1, \dots, e_n)} \cdot \widetilde{M}_{(e_1, \dots, e_n)}(z)$$

where $\widetilde{M}_{(e_1, \dots, e_n)}(z)$ is the truncation of $M_{(e_1, \dots, e_n)}(z)$ by $z^{n-2} = 0$: When the Euler class of the obstruction bundle for (e_1, \dots, e_n) is given as a polynomial $P(\psi_n)$ in ψ_n , then the contribution to the J -function is the ψ_n^{n-3} -coefficient of $P(\psi_n) \frac{1}{z - \psi_n}$. This coefficient is given by z^{-n+2} times the truncation of $P(z)$.

Independently of n , this shows that $J^{X; \text{weighted}}$ is obtained from I^{tw} by removing all terms of non-negative degree in z except the first two; such terms would correspond geometrically to a negative number of ψ_n -insertions.

4. WEIGHT CHANGE

The goals of this section are the wall-crossing theorems 4.3.1 and 4.3.2.

⁶The formula on page 9 ibid. defining the “modification factor” $M_\theta(z)$ has to be applied with s_i specialized such that $e^{s(ch)}$ for the Chern character ch of some bundle E gives the Euler class of $-[E]$.

⁷It has the same image as the twisted J -function, namely Givental's Lagrangian cone \mathcal{L}_X .

4.1. Preparations. We begin with several lemmata we will need in the proof.

Lemma 4.1.1. *Let D_1, D_2 be two smooth divisors with transversal intersection on a smooth Deligne-Mumford stack X . Let \tilde{X} be the blow-up of X at their intersection, with exceptional divisor E and proper transforms \tilde{D}_1 and \tilde{D}_2 . On the other hand, consider the r -th root construction $X_{(D_1,r),(D_2,r)}$ and its blowup Z at the intersection of the two gerbes \mathcal{D}_1 and \mathcal{D}_2 lying over D_1 and D_2 , respectively. Then Z is isomorphic to the r -th root construction $\tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)}$.*

Proof. Let $\tilde{\mathcal{D}}_i$ and \mathcal{E} be the gerbes in $\tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)}$ lying over \tilde{D}_i and E , respectively. The line bundles $\mathcal{O}(\tilde{\mathcal{D}}_i \otimes \mathcal{E})$ with their canonical sections are r -th roots of the pull-backs of $D_i \subset X$, determining a morphism

$$f: \tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)} \rightarrow X_{(D_1,r),(D_2,r)}.$$

The pull-back of the ideal sheaf of the intersection $\mathcal{D}_1 \cap \mathcal{D}_2$ along f is the ideal sheaf of \mathcal{E} : this is easy to see locally, where we can assume that \mathcal{D}_i is cut out by an equation $(s_i = 0)$; its pull-back f^*s_i cuts out $\mathcal{E} \cup \tilde{\mathcal{D}}_i$. Since \mathcal{E} is Cartier, the universal property of blow-ups yields a map

$$g: \tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)} \rightarrow Z.$$

To go the other way, we first show that $Z \rightarrow X$ lifts to \tilde{X} . The preimage of $D_1 \cap D_2$ in $X_{(D_1,r),(D_2,r)}$ is $\mathcal{D}_1^r \cap \mathcal{D}_2^r$. One can check after étale base change to a scheme that the preimage of this in Z is r times the exceptional divisor, hence is Cartier. So the universal property of blowups gives us a morphism $Z \rightarrow \tilde{X}$. The preimage of E under this morphism is r times the exceptional divisor of Z , and it follows that the preimage of \tilde{D}_i is r times the proper transform of \mathcal{D}_i . This gives us a lifting to

$$h: Z \rightarrow \tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)}.$$

As neither Z nor $\tilde{X}_{(\tilde{D}_1,r),(\tilde{D}_2,r),(E,r)}$ have nontrivial automorphisms over the identity of X , both gh and hg must be (2-isomorphic to) the identity. \square

Lemma 4.1.2. *Let X, Y be Deligne-Mumford stacks, and let $f: X \rightarrow Y$ be a composition of r -th root constructions and blow-ups at regularly embedded centers (i.e. the normal sheaf of the center is a vector bundle). Then $\mathbb{R}f_*\mathbb{L}f^*\mathcal{F} = \mathcal{F}$ for any quasi-coherent sheaf \mathcal{F} on Y .*

Proof. By the projection formula, it is enough to prove $\mathbb{R}f_*\mathcal{O}_X = \mathcal{O}_Y$. For r -th root constructions, it is obvious that the higher direct images vanish, and by [Cad07] $f_*\mathcal{O}_X = \mathcal{O}_Y$. For blow-ups, this is well-known in the case of schemes. Since blow-ups are representable, one can reduce to the case of schemes by taking an étale base change to a scheme covering Y . \square

Lemma 4.1.3. (*Base change*) *Let*

$$\begin{array}{ccc} Y' & \xrightarrow{h} & Y \\ f' \downarrow & \square & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

be a 2-cartesian square of Deligne-Mumford stacks where X is quasi-compact, f is quasi-compact and quasi-separated, and Y and X' are tor-independent over X . Let $\mathcal{E} \in D^b(Y)$ be a complex having quasi-coherent cohomology. Suppose either that g has finite tor-dimension or that \mathcal{E} has flat, finite amplitude relative to f . Then there is a natural isomorphism

$$\mathbb{L}g^*\mathbb{R}f_*\mathcal{E} \rightarrow \mathbb{R}f'_*\mathbb{L}h^*\mathcal{E}.$$

We will apply this in the case where f is flat and \mathcal{E} is a line bundle on Y .

Proof. The existence of a natural morphism $\mathbb{L}g^*\mathbb{R}f_*\mathcal{E} \rightarrow \mathbb{R}f'_*\mathbb{L}h^*\mathcal{E}$ follows from the adjointness of pullback and pushforward. Indeed, the natural morphism $\mathcal{E} \rightarrow \mathbb{R}h_*\mathbb{L}h^*\mathcal{E}$ determines a morphism

$$\mathbb{R}f_*\mathcal{E} \rightarrow \mathbb{R}(fh)_*\mathbb{L}h^*\mathcal{E} = \mathbb{R}(gf')_*\mathbb{L}h^*\mathcal{E},$$

which is equivalent to the morphism above by adjointness of $\mathbb{R}g_*$ and $\mathbb{L}g^*$.

For schemes, the proposition is the same as [SGA71, IV, 3.1.0], and the reduction to the case of schemes is identical to the proof of [LMB00, 13.1.9]. \square

Lemma 4.1.4. *Let $\tau: \tilde{X} \rightarrow X$ be the blow-up of a smooth Deligne-Mumford stack X at a smooth center $Z \subset X$ of codimension two, with normal bundle N and exceptional divisor E . Then for $n \geq 0$*

$$[R^1\tau_*\mathcal{O}(n \cdot E)] = \sum_{k=0}^{n-2} [\Lambda^2 N \otimes \text{Sym}^k N].$$

Proof. As a blow-up is representable, it is sufficient to check this for schemes. Due to the short exact sequences $\mathcal{O}((n-1) \cdot E) \rightarrow \mathcal{O}(nE) \rightarrow \mathcal{O}_E(-n)$ on \tilde{X} , it follows from induction if we show $R^1\pi_*\mathcal{O}_E(-n) = [\Lambda^2 N \otimes \text{Sym}^{n-2} N]$. This is easily checked by Serre duality, since

$$R^0\pi_*\mathcal{O}_E(n) = \text{Sym}^n(N^\vee)$$

and the relative dualizing sheaf of $\tau|_E$ is $\mathcal{O}_E(-2) \otimes \tau^*(\Lambda^2 N)^{-1}$. \square

4.2. Constructing the reduction map. Consider two weight data $\mathcal{A} > \mathcal{B}$; then there is a reduction morphism $\rho_{\mathcal{B},\mathcal{A}}: \overline{M}_{0,\mathcal{A}} \rightarrow \overline{M}_{0,\mathcal{B}}$. Further, we assume the following property:

- (*) There is exactly one 2-partition $\sigma = (T_0, T_0^C)$ of $[n]$ such that σ is \mathcal{A} -stable but \mathcal{B} -unstable.

In other words, there is just one wall between \mathcal{A} and \mathcal{B} in the chamber decomposition of the set of weight data discussed in section 3.1. Specifically, this means that $\sum_{i \in T_0} a_i > 1$ but $\sum_{i \in T_0} b_i \leq 1$, and that T_0 is the only such subset of $[n]$.

Then $\rho_{\mathcal{B},\mathcal{A}}: \overline{M}_{0,\mathcal{A}} \rightarrow \overline{M}_{0,\mathcal{B}}$ is the blow-up of $\overline{M}_{0,\mathcal{B}}$ at the locus Z_{T_0} of curves where all $s_i, i \in T_0$ agree; the exceptional divisor is D^{T_0} . The universal curve $\overline{C}_{\mathcal{A}}$ is obtained from $\overline{C}_{\mathcal{B}}$ in two steps:

- (1) Blowing up at the preimage $\pi^{-1}Z_{T_0}$ of Z_{T_0} , i.e. taking the base change for $\rho_{\mathcal{B},\mathcal{A}}$; we denote the resulting family over $\overline{M}_{0,\mathcal{A}}$ by $\overline{C}_{\mathcal{B}/\mathcal{A}}$.
- (2) Blowing up at the preimage (with respect to the previous blow-up) of the common image $s_i(Z_{T_0})$ for any $i \in T_0$. (See [BM06, Remark 3.1.2].)

There is a canonical map

$$\rho_{\mathcal{B},\mathcal{A}}^{(1)}: \overline{M}_{0,\mathcal{A}}^{(1)} \rightarrow \overline{M}_{0,\mathcal{B}}^{(1)}.$$

This follows from the fact that the pull-back of any boundary divisor $D^S \subset \overline{M}_{0,\mathcal{B}}$ is just the corresponding divisor $D^S \subset \overline{M}_{0,\mathcal{A}}$, and that the r -th root construction commutes with such base change.

A matching map $\overline{C}_{\mathcal{A}}^{(1)} \rightarrow \overline{C}_{\mathcal{B}}^{(1)}$ does not exist in general, for the following reason: the preimage of a section $s_i \subset \overline{C}_{\mathcal{B}}$ with $i \in T_0$ is the union of the corresponding section $s_i \in \overline{C}_{\mathcal{A}}$ with the exceptional divisor $D^{T_0, n+1}$ of the second blow-up step in the construction of $\overline{C}_{0,\mathcal{A}}$ above. However, for example when $\prod_{i \in T_0} e_i = 1$, there is no root construction along the divisor $D^{T_0, n+1}$ in the construction of $\overline{C}_{\mathcal{A}}^{(1)}$ at all, and so an r_i -th root of the pull-back of $s_i \subset \overline{C}_{0,\mathcal{B}}$ does not exist.

In order to compare the Hodge bundles, we will later construct some auxiliary spaces to overcome this problem.

For simplicity, we make the following additional assumption:

(**) We assume that for T_0 as in (*), we have $n \notin T_0$.

(This simplifies the computation with respect to our choice of $L_{1,\mathcal{A}}$ in definition 3.3.1 and, on the other hand, always holds when we start with the \mathbb{P}^{n-3} -weight data used in section 3.4.)

Lemma 4.2.1. *Assuming (*) and (**), we have $\rho_{\mathcal{B},\mathcal{A}}^* L_{2,\mathcal{B}} = L_{2,\mathcal{A}}$*

This is immediate from $L_{2,\mathcal{A}} = s_n^*(L_{1,\mathcal{A}})^r$, as s_n does not meet any of the divisors appearing in the definition (3.3.1) of $L_{1,\mathcal{A}}$, except itself. As a consequence, it follows that:

Corollary 4.2.2. *There is a well-defined reduction map*

$$\rho_{\mathcal{B},\mathcal{A}}: \overline{M}_{0,\mathcal{A},\mathcal{E}}^{\mu_r} \rightarrow \overline{M}_{0,\mathcal{B},\mathcal{E}}^{\mu_r}.$$

4.3. Weight change and Hodge bundles. We continue with the assumptions (*) and (**) from the previous section. We want to compare $\rho_{\mathcal{B},\mathcal{A}}^* H_{\mathcal{B}}^w$ and $H_{\mathcal{A}}^w$ in the K -group of $\overline{M}_{0,\mathcal{A},\mathcal{E}}^{\mu_r}$, where due to projection formula $H_{\mathcal{A}}^w$ can be computed by

$$H_{\mathcal{A}}^w = R^1 \pi_* L_{1,\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}}.$$

We will introduce several auxiliary spaces; the goal is to have a reduction map as a smooth blow-up between spaces that are very close to the universal curves. This is achieved in the map τ below.

Let $A_{\mathcal{A}}$ be the intermediate space obtained from $\overline{C}_{0,\mathcal{A}}^{\mu_r}$ by forgetting the root construction along all sections. In other words, it is constructed from the universal curve $\overline{C}_{\mathcal{A}}$ by the r_T -th root construction for every divisor D^T where $T \subset [n+1]$ and T is $\mathcal{A} \cup \{0\}$ -stable, followed by the base change along $\overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r} \rightarrow \overline{M}_{0,\mathcal{A}}^{(1)}$. Let $\nu_{\mathcal{A}}: \overline{C}_{0,\mathcal{A}}^{\mu_r} \rightarrow A_{\mathcal{A}}$ be the induced map, and $\pi'_{\mathcal{A}}: A_{\mathcal{A}} \rightarrow \overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r}$ the projection to the moduli space.

Then

$$(4.3.1) \quad \nu_{\mathcal{A}*}(L_{1,\mathcal{A}}^w) = \mathcal{O} \left(\sum_{i=1}^n [wd_i] s_i + \sum_{\substack{T \subset [n-1] \\ (T, T^C) \text{ is } \mathcal{A}\text{-stable}}} wd_T \cdot D^{T,n+1} \right)$$

Now let $A_{\mathcal{B}/\mathcal{A}}$ be the base change of $A_{\mathcal{B}} \rightarrow \overline{M}_{0,\mathcal{B},\varepsilon}^{\mu_r}$ along $\rho_{\mathcal{B},\mathcal{A}}$. While a map $A_{\mathcal{A}} \rightarrow A_{\mathcal{B}/\mathcal{A}}$ exists, we prefer not to use it and instead consider two more additional spaces: Pick any section s_{j_0} with $j_0 \in T_0$ and let $A'_{\mathcal{A}} = (A_{\mathcal{A}})_{s_{j_0}, r_T}$ be the stack obtained from $A_{\mathcal{A}}$ by adding the r_T -th root construction at the j_0 -th section. We define $A'_{\mathcal{B}/\mathcal{A}}$ analogously.

$$\begin{array}{ccccc} \overline{C}_{0,\mathcal{A}}^{\mu_r} & & A'_{\mathcal{A}} & \xrightarrow{\tau} & A'_{\mathcal{B}/\mathcal{A}} \\ & \searrow \nu_{\mathcal{A}} & \downarrow \epsilon_{\mathcal{A}} & & \downarrow \epsilon_{\mathcal{B}/\mathcal{A}} \\ & & A_{\mathcal{A}} & & A_{\mathcal{B}/\mathcal{A}} \xrightarrow{\rho_{\mathcal{B},\mathcal{A}}} A_{\mathcal{B}} \\ & & \searrow \pi'_{\mathcal{A}} & & \downarrow \pi'_{\mathcal{B}} \\ & & & & \overline{M}_{0,\mathcal{A},\varepsilon}^{\mu_r} \xrightarrow{\rho_{\mathcal{B},\mathcal{A}}} \overline{M}_{0,\mathcal{B},\varepsilon}^{\mu_r} \end{array}$$

Applying lemma 4.1.1 to the divisors s_{j_0} and $\pi^{-1}D^T$ on the coarse moduli space of $A_{\mathcal{B}/\mathcal{A}}$, we see that the map τ is the blow-up at the intersection of the tautological gerbes over s_{j_0} and $(\pi'_{\mathcal{B}/\mathcal{A}})^{-1}D^{T_0}$ in $A'_{\mathcal{B}/\mathcal{A}}$.

Setting $L_{1,\mathcal{B}/\mathcal{A}}^w = \rho_{\mathcal{B},\mathcal{A}}^* \nu_{\mathcal{B}*} L_{1,\mathcal{B}}^w$, we have

$$\begin{aligned} (\rho_{\mathcal{B},\mathcal{A}})^* H_{\mathcal{B}}^w &= (\rho_{\mathcal{B},\mathcal{A}})^* R^1(\pi'_{\mathcal{B}})_* \nu_{\mathcal{B}*} L_{1,\mathcal{B}}^w \otimes L_{2,\mathcal{B}}^{-\frac{w}{r}} \\ &= R^1(\pi'_{\mathcal{B}/\mathcal{A}})_* \rho_{\mathcal{B},\mathcal{A}}^* \nu_{\mathcal{B}*} L_{1,\mathcal{B}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}} \quad (\text{lemma 4.1.3}) \\ (4.3.2) \quad &= R^1(\pi'_{\mathcal{B}/\mathcal{A}})_* L_{1,\mathcal{B}/\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}} \\ &= R^1(\pi'_{\mathcal{B}/\mathcal{A}} \epsilon_{\mathcal{B}/\mathcal{A}} \tau)_* (\epsilon_{\mathcal{B}/\mathcal{A}} \tau)^* L_{1,\mathcal{B}/\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}} \quad (\text{lemma 4.1.2}) \end{aligned}$$

on the other hand,

$$\begin{aligned} H_{\mathcal{A}}^w &= R^1(\pi'_{\mathcal{A}})_* \nu_{\mathcal{A}*} L_{1,\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}} \\ &= R^1(\pi'_{\mathcal{A}} \epsilon_{\mathcal{A}})_* \epsilon_{\mathcal{A}}^* \nu_{\mathcal{A}*} L_{1,\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}} \quad (\text{lemma 4.1.2}) \\ (4.3.3) \quad &= R^1(\pi'_{\mathcal{B}/\mathcal{A}} \epsilon_{\mathcal{B}/\mathcal{A}} \tau)_* \epsilon_{\mathcal{A}}^* \nu_{\mathcal{A}*} L_{1,\mathcal{A}}^w \otimes (L_{2,\mathcal{A}})^{-\frac{w}{r}}. \end{aligned}$$

Now for any $T \subset [n-1]$ such that (T, T^C) is \mathcal{B} -stable, it is also \mathcal{A} -stable, and

$$(\rho_{\mathcal{B}, \mathcal{A}} \epsilon_{\mathcal{B}/\mathcal{A}} \tau)^*(D^{T, n+1}) = \epsilon_{\mathcal{A}}^*(D^{T, n+1}).$$

The pull-back of the divisor class of sections is given by

$$(\rho_{\mathcal{B}, \mathcal{A}} \epsilon_{\mathcal{B}/\mathcal{A}} \tau)^*(s_i) = \begin{cases} s_i & \text{if } i \notin T_0 \\ s_i + D^{T_0, n+1} & \text{if } i \in T_0 \end{cases}$$

Using these formulae and equation (4.3.1) for \mathcal{A} and \mathcal{B} , respectively, yields

$$(\epsilon_{\mathcal{A}})^* \nu_{\mathcal{A}*} L_{1, \mathcal{A}}^w = (\epsilon_{\mathcal{B}/\mathcal{A}} \tau)^* L_{1, \mathcal{B}/\mathcal{A}}^w \otimes \mathcal{O} \left(\sum_{i \in T_0} (wd_i - \lfloor wd_i \rfloor) \cdot D^{T_0, n+1} \right).$$

Note that $\delta_{T_0}^w = \sum_{i \in T_0} (wd_i - \lfloor wd_i \rfloor)$. The projection formula yields

$$(4.3.4) \quad \mathbb{R}(\epsilon_{\mathcal{B}/\mathcal{A}} \tau)_*(\epsilon_{\mathcal{A}})^* \nu_{\mathcal{A}*} L_{1, \mathcal{A}}^w = L_{1, \mathcal{B}/\mathcal{A}}^w \otimes \mathbb{R}(\epsilon_{\mathcal{B}/\mathcal{A}} \tau)_* \mathcal{O}(\delta_{T_0}^w \cdot D^{T_0, n+1}).$$

Combining equations (4.3.2, 4.3.3, 4.3.4), and using $R^0(\epsilon_{\mathcal{B}/\mathcal{A}} \tau)_* \mathcal{O}(\delta_{T_0}^w \cdot D^{T_0, n+1}) = \mathcal{O}$, it follows that

$$(4.3.5) \quad [H_{\mathcal{A}}^w] = [\rho_{\mathcal{B}, \mathcal{A}}^* H_{\mathcal{B}}^w] + \left[(\pi'_{\mathcal{B}/\mathcal{A}})_* \left(R^1(\epsilon_{\mathcal{B}/\mathcal{A}} \tau)_* \mathcal{O}(\delta_{T_0}^w D^{T_0, n+1}) \otimes L_{1, \mathcal{B}/\mathcal{A}}^w \right) \otimes L_{2, \mathcal{A}}^{-\frac{w}{r}} \right]$$

Write $\delta_{T_0}^w$ as the fraction $\frac{p_0}{r_0}$, where $r_0 = r_{T_0}$. The exceptional divisor of τ is $\mathcal{D}^{T_0, n+1} = \frac{1}{r_0} D^{T_0, n+1}$; so by lemma 4.1.4

$$(4.3.6) \quad [R^1 \tau_* \mathcal{O}(\delta_{T_0}^w \cdot D^{T_0, n+1})] = \sum_{k=0}^{p_0-2} [\Lambda^2 N \otimes \text{Sym}^k N],$$

where N is the normal bundle to the center of the blow-up τ . To compute the right-hand side, we introduce additional normal bundles. Let $N_{\mathfrak{s}_{j_0}}$ be the normal bundle of the gerbe \mathfrak{s}_{j_0} over s_{j_0} in $A'_{\mathcal{B}/\mathcal{A}}$, let $N_{\mathcal{D}^{T_0}}$ be the normal bundle to the gerbe \mathcal{D}^{T_0} over D^{T_0} in $\overline{M}_{0, \mathcal{A}, \mathcal{E}}^{\mu_r}$, and let $N_{s_{j_0}}$ be the normal bundle to s_{j_0} in $A_{\mathcal{B}/\mathcal{A}}$ (equivalently, $N_{s_{j_0}}$ is the relative tangent bundle of $\pi'_{\mathcal{B}/\mathcal{A}}$ restricted to s_{j_0}). Then $N_{\mathfrak{s}_{j_0}}$ is the restriction of the tautological bundle of the r_0 -th root construction $\epsilon_{\mathcal{B}/\mathcal{A}}$ at the section s_{j_0} to the gerbe, and

$$(4.3.7) \quad (\epsilon_{\mathcal{B}/\mathcal{A}})_* N_{\mathfrak{s}_{j_0}}^k = \begin{cases} N_{s_{j_0}}^{\frac{k}{r_0}} & \text{if } r_0 \text{ divides } k, \\ 0 & \text{otherwise.} \end{cases}$$

The section s_{j_0} induces a splitting of the tangent bundle of $A_{\mathcal{B}/\mathcal{A}}$ along s_{j_0} into the relative tangent bundle and the push-forward of the tangent bundle of $\overline{M}_{0, \mathcal{A}, \mathcal{E}}^{\mu_r}$ along s_{j_0} . This induces a splitting of N as

$$N = \left((\epsilon_{\mathcal{B}/\mathcal{A}} \pi'_{\mathcal{B}/\mathcal{A}})^* N_{\mathcal{D}^{T_0}} \oplus N_{\mathfrak{s}_{j_0}} \right) \Big|_{Z_{T_0}}$$

Applying this splitting to equation (4.3.6), we obtain

$$[R^1\tau_*\mathcal{O}(\delta_{T_0}^w \cdot D^{T_0,n+1})] = \sum_{k=0}^{p_0-2} \sum_{a+b=k} [N_{s_{j_0}}^{b+1} \otimes (\epsilon_{\mathcal{B}/\mathcal{A}}\pi'_{\mathcal{B}/\mathcal{A}})^* N_{\mathcal{D}^{T_0}}^{a+1}].$$

We can rewrite the summation as $\sum_{b=0}^{p_0-2} \sum_{a=0}^{p_0-b-2}$ or, equivalently, $\sum_{b=0}^{p_0-1} \sum_{a=0}^{p_0-b-2}$. By equation (4.3.7), only the terms with $mr_0 = b+1$ for some $m \in \mathbb{Z}$ are surviving the push-forward along $\epsilon_{\mathcal{B}/\mathcal{A}}$, which yields:

$$[R^1(\epsilon_{\mathcal{B}/\mathcal{A}}\tau)_*\mathcal{O}(\delta_{T_0}^w \cdot D^{T_0,n+1})] = \sum_{m=1}^{\lceil \delta_{T_0}^w \rceil - 1} \sum_{a=0}^{p_0-r_0m-1} [N_{s_{j_0}}^m \otimes (\pi'_{\mathcal{B}/\mathcal{A}})^* N_{\mathcal{D}^{T_0}}^{a+1}]$$

Combining this with equation (4.3.5) yields the wall-crossing theorem in the K -group:

Theorem 4.3.1. *The generalized dual Hodge bundles $H_{\mathcal{A}}^w$ and $H_{\mathcal{B}}^w$ can be related in the K -group of $\overline{M}_{0,\mathcal{A},\mathcal{E}}^{\mu_r}$ as follows:*

$$(4.3.8) \quad [H_{\mathcal{A}}^w] = [\rho_{\mathcal{B},\mathcal{A}}^* H_{\mathcal{B}}^w] + \sum_{m=1}^{\lceil \delta_{T_0}^w \rceil - 1} \sum_{a=1}^{p_0-r_0m} [N_{\mathcal{D}^{T_0}}^a \otimes s_{j_0}^* N_{s_{j_0}}^m \otimes s_{j_0}^* L_{1,\mathcal{B}/\mathcal{A}}^w \otimes L_{2,\mathcal{A}}^{-\frac{w}{r}}]$$

where s_{j_0} is the section $s_{j_0}: \overline{M}_{0,\mathcal{A},\mathcal{E}}^{\mu_r} \rightarrow A_{\mathcal{B}/\mathcal{A}}$.

Let α be the first Chern class of the line bundle $s_{j_0}^* \nu_{\mathcal{B}}^* L_{1,\mathcal{B}}^w \otimes L_{2,\mathcal{B}}^{-\frac{w}{r}}$ on $\overline{M}_{0,\mathcal{B},\mathcal{E}}^{\mu_r}$. The first Chern class of $s_{j_0}^* N_{s_{j_0}}$ on $\overline{M}_{0,\mathcal{B},\mathcal{E}}^{\mu_r}$ is $-\psi_{j_0}$, and so the first Chern class of $\mathcal{K} := s_{j_0}^* N_{s_{j_0}}^m \otimes s_{j_0}^* L_{1,\mathcal{B}/\mathcal{A}}^w \otimes L_{2,\mathcal{A}}^{-\frac{2}{r}}$ is $\rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})$.

Tensoring the short exact sequences

$$0 \rightarrow \mathcal{O}(\frac{1}{r_0}D^{T_0})^{a-1} \rightarrow \mathcal{O}(\frac{1}{r_0}D^{T_0})^a \rightarrow N_{\mathcal{D}^{T_0}}^a \rightarrow 0$$

with \mathcal{K} yields Chern classes for all summands of the right hand side of equation (4.3.8), and thus the following formula relating the Chern classes of the dual Hodge bundles:

$$\begin{aligned} c(H_{\mathcal{A}}^w) &= \rho_{\mathcal{B},\mathcal{A}}^*(c(H_{\mathcal{B}}^w)) \cdot \prod_{m=1}^{\lceil \delta_{T_0}^w \rceil - 1} \prod_{a=1}^{p_0-r_0m} \frac{1 + \frac{a}{r_0}D^{T_0} + \rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})}{1 + \frac{a-1}{r_0}D^{T_0} + \rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})} \\ &= \rho_{\mathcal{B},\mathcal{A}}^*(c(H_{\mathcal{B}}^w)) \cdot \prod_{m=1}^{\lceil \delta_{T_0}^w \rceil - 1} \frac{1 + (\delta_{T_0}^w - m)D^{T_0} + \rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})}{1 + \rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})} \\ &= \rho_{\mathcal{B},\mathcal{A}}^*(c(H_{\mathcal{B}}^w)) \cdot \prod_{m=1}^{\lceil \delta_{T_0}^w \rceil - 1} \left(1 + \frac{(\delta_{T_0}^w - m)D^{T_0}}{1 + \rho_{\mathcal{B},\mathcal{A}}^*(\alpha - m\psi_{j_0})} \right) \end{aligned}$$

Equation (4.3.8) implies that this formula does not depend on α itself but only on its restriction $\alpha \cdot Z_{T_0}$ to the center of the blowup-part of $\rho_{\mathcal{B},\mathcal{A}}$. The Chern class

of $L_{1,\mathcal{B}}$ on $\overline{C}_{0,\mathcal{B}}^{\mu_r}$ is by construction equal to the Chern class of $\pi_{\mathcal{B}}^* L_{2,\mathcal{B}}^{\frac{1}{r}}$. Hence α can be computed as the difference of the Chern classes of $\nu_{\mathcal{B}*} L_{1,\mathcal{B}}^w$ and $L_{1,\mathcal{B}}^w$, which is given by $\sum_{i=1}^n ([wd_i] - wd_i) s_i = -\sum_{i=1}^n \delta_i^w s_i$. Pulling this back via s_{j_0} , and ignoring everything that restricts as zero to Z_{T_0} , gives $\alpha = -\sum_{i \in T_0} \delta_i^w (-\psi_{j_0}) = \delta_{T_0}^w \psi_{j_0}$.

We claim that the restriction of $\rho_{\mathcal{B},\mathcal{A}}^* \psi_{j_0}$ to D^{T_0} is the ψ -class of the node on the component corresponding to the complement of T_0 . To see this, note that the irreducible component over D^{T_0} corresponding to T_0^C is the pull-back of the family over Z_{T_0} , with s_{j_0} being pulled back to the node.

Since in the final formula, after expanding the fraction using a geometric series, $\rho_{\mathcal{B},\mathcal{A}}^* \psi_{j_0}$ only appears in monomials that also have a factor of D^{T_0} , we can replace $\rho_{\mathcal{B},\mathcal{A}}^* \psi_{j_0}$ by ψ_{T_0} ; here ψ_{T_0} is for now any divisor on that restricts as the ψ -class of the node to D^{T_0} (but see (A.1.4) for a somewhat canonical global definition). So the formula simplifies further:

Theorem 4.3.2. *Assume that there is a single wall w_{T_0} as defined in (3.1.1) between the two weight data $\mathcal{A} \geq \mathcal{B}$. The Chern classes of the generalized dual Hodge bundles can be related on $\overline{M}_{0,\mathcal{A},\mathcal{E}}^{\mu_r}$ as follows:*

$$c(H_{\mathcal{A}}^w) = \rho_{\mathcal{B},\mathcal{A}}^*(c(H_{\mathcal{B}}^w)) \cdot \prod_{p=\langle \delta_{T_0}^w \rangle}^{\delta_{T_0}^w - 1} \left(1 + \frac{pD^{T_0}}{1 + p\psi_{T_0}} \right)$$

5. CHERN CLASS FORMULA

5.1. Main theorem. Theorem 4.3.2 and corollary 3.4.2 immediately give a closed formula for the equivariant Euler class of the generalized dual Hodge bundle.

Assume that μ_r is acting diagonally on \mathbb{C}^N with weights w_1, \dots, w_N . Given $e_1, \dots, e_n \in \mu_r$, let $\delta_i^{(a)} \in [0, 1)$ be the age of e_i acting on the a -th coordinate direction, i.e. $e^{2\pi i \delta_i^{(a)}} = e_i^{w_a}$. For all subsets $T \subset [n]$, let $\delta_T^{(a)} = \sum_{i \in T} \delta_i^{(a)}$.

Theorem 5.1.1. *On the connected component $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ of the moduli space of twisted stable maps $\overline{M}_{0,n}(B\mu_r)$, the equivariant Euler class of the obstruction bundle is given as*

$$e_T([R^1\pi_* f^* \mathbb{C}^N]) = \prod_{a=1}^N \prod_{p=\langle \delta_{[n-1]}^{(a)} \rangle}^{\delta_{[n-1]}^{(a)} - 1} (t_a - p\psi_n) \cdot \prod_{\substack{T \subset [n-1] \\ 2 \leq |T|}} \prod_{p=\langle \delta_T^{(a)} \rangle}^{\delta_T^{(a)} - 1} \left(1 + \frac{pD^T}{t_a + p\psi_T} \right)$$

Proof. If we start with weight data $\mathcal{A} = (\frac{1}{n-2}, \dots, \frac{1}{n-2}, 1)$ as in section 3.4, we can choose a path in $[0, 1]^n$ leading to $\mathcal{A} = (1, 1, \dots, 1)$ such that we pass every wall $w_T = \{a_i \mid \sum_{i \in T} a_i = 1\}$ for $T \subsetneq [n-1]$, $|T| \geq 2$ exactly once, and only one wall at a time. By theorem 4.3.2, we pick up exactly the factor in the above product corresponding to D^T when we cross the wall w_T , after we set $t_a = 1$. To get the equivariant Euler class from the total Chern class, we just have to multiply the i -th

Chern class of the higher direct image of the a -th coordinate direction $\mathbb{C} \subset \mathbb{C}^N$ with t_a^{k-i} , as the torus is acting trivially on the moduli space, and linearly with multiplication by t_a on the fibers of the vector bundle. \square

5.2. Remarks on the formula. Using the notations $D^{[n-1]} = -\psi_n$, $D^i = -\psi_i$ and $\psi_{[n-1]} = 0$ as explained in the appendix, the formula can be written in more compact form:

(5.2.1)

$$e_T([R^1\pi_*f^*\mathbb{C}^N] - [R^0\pi_*f^*\mathbb{C}^N]) = \prod_{a=1}^N t_a^{\delta_{[n]}^{(a)}-1} \prod_{\emptyset \neq T \subseteq [n-1]} \prod_{p=\langle \delta_T^{(a)} \rangle}^{\delta_T^{(a)}-1} \frac{t_a + p\psi_T + pD^T}{t_a + p\psi_T}$$

Here $\delta_{[n]}^{(a)} - 1$ is the virtual dimension of the contribution from the a -th coordinate direction $\mathbb{C} \subset \mathbb{C}^N$ to obstruction bundle.

The convenience of this formulation is that it remains correct (up to an overall power of t_a) as long as $\delta_{\{i\}}^{(a)}$ is any real number such that $e_i^{w_a} = e^{2\pi i \delta_{\{i\}}^{(a)}}$ (if we still define $\delta_T^{(a)} = \sum_{i \in T} \delta_{\{i\}}^{(a)}$). This is shown in the appendix, see lemma A.1.1.

This version of the formula also gives the correct answer for the necessary localization computation in case all e_i act trivially on one of the coordinate direction, that is if $e_i^{w_a} = 1$ for some a and all i . In that case, $\overline{M}_{0,n}(e_1, \dots, e_n; B\mu_r)$ is the fixed point locus of $\overline{M}_{0,n}(e_1, \dots, e_n; [\mathbb{C}^N/\mu_r])$ (instead of being isomorphic to it). The factor of $\frac{1}{t_a}$ we get in the above formula is the contribution of the a -th coordinate direction to the inverse of the equivariant Euler class of the normal bundle of the fixed point locus.

6. RECURSIONS FOR GROMOV-WITTEN INVARIANTS

6.1. Inclusion-exclusion principle. The formula gives particularly nice recursions when the invariants are (almost) non-equivariant. To expand the formula, we use the following fact, which we think of as a generalized inclusion-exclusion principle:

Lemma 6.1.1. *Let S be a partially ordered set. Let $U(S)$ be the set of non-empty subsets $I \subset S$ such that no two elements of I are comparable. For every subset $I \subset S$, let $C(I) \subset S$ be the “ordered complement” of I : the set of elements of S that are not less than or equal to any element of I . Then:*

$$(6.1.1) \quad \prod_{T \in S} (1 + x_T) = 1 + \sum_{I \in U(S)} (-1)^{|I|+1} \prod_{T \in I} x_T \prod_{T \in C(I)} (1 + x_T)$$

Proof. For any subset $J \subset S$, the monomial $\prod_{T \in J} x_T$ appears in the right-hand side of the above product whenever I is a subset of the set of minimal elements of J . It is easily checked that it overall has coefficient one. \square

We use this for S being the set of subsets $T \subset [n-1]$ with $2 \leq |T| \leq n-2$, ordered by inclusion, and

$$\begin{aligned} x_T &= -1 + \prod_{a=1}^N \prod_{p=\langle \delta_T^{(a)} \rangle}^{\delta_T^{(a)}-1} \left(1 + \frac{pD^T}{t_a + p\psi_T} \right) \\ &= \frac{\prod_{a,p}(1 - t_a^{-1}p\psi_{TC}) - \prod_{a,p}(1 + t_a^{-1}p\psi_T)}{\prod_{a,p}(1 + t_a^{-1}p\psi_T)}, \end{aligned}$$

where $\psi_{TC} = -D^T - \psi_T$ (see Appendix). Then x_T is a class with support on D^T . Thus, if T, T' are not comparable, then $x_T x_{T'}$ can only be non-zero if T and T' are disjoint; hence the expansion of lemma 6.1.1 reduces to a sum over a combination of pairwise disjoint subsets $T_1, \dots, T_k \subset [n-1]$. Since every T_i has size at least 2, the datum of $\{T_1, \dots, T_k\}$ can be identified with a partition \mathcal{P} of $[n-1]$; given \mathcal{P} , we write $\mathcal{P}_{\geq 2}$ for the sets in \mathcal{P} that have size at least 2, recovering the list of the T_i . We can similarly simplify the second product of equation (6.1.1) to a product over T which are either disjoint from or fully contain T_i , for all i ; in other words, we can identify T with a subset of the quotient set $[n-1]/(\mathcal{P})$ having at least 2 elements. Thus:

$$(6.1.2) \quad e_T(R^1\pi_* f^* \mathbb{C}^N) = \left(1 + \sum_{\mathcal{P}} (-1)^{|\mathcal{P}_{\geq 2}|+1} \prod_{T \in \mathcal{P}_{\geq 2}} x_T \prod_{\substack{T \subset [n-1]/(\mathcal{P}) \\ 2 \leq |T|}} (1 + x_T) \right) \prod_{a=1}^N \prod_{p=\langle \delta_{[n-1]}^{(a)} \rangle}^{\delta_{[n-1]}^{(a)}-1} (t_a - p\psi_n)$$

where the sum goes over all non-trivial partitions \mathcal{P} of $[n-1]$ (excluding the partitions of size 1 and $n-1$), and we identify a subset $T \subset [n-1]/(\mathcal{P})$ with its preimage in $[n-1]$.

The class associated to the partition \mathcal{P} in the above expansion has support on $\bigcap_{T \in \mathcal{P}_{\geq 2}} D^T$, which explains why we call it an inclusion-exclusion principle. This intersection is a moduli space of comb curves as in the figure on page 29.

6.2. Non-equivariant recursions for $[\mathbb{C}^3/\mu_r]$. Let μ_r act non-trivially on \mathbb{C}^3 so that it leaves the volume form of \mathbb{C}^3 invariant. Up to isomorphism of μ_r , we may assume that the generator is acting with age 1; then the weights w_1, w_2, w_3 of the one-dimensional representations satisfy $w_1 + w_2 + w_3 = r$. The age of the action of a non-trivial group element e_i is given by $\text{age}(e_i, \mathbb{C}^3) = \text{age}(e_i^{w_1}) + \text{age}(e_i^{w_2}) + \text{age}(e_i^{w_3})$. In this section we will develop recursions for invariants of the form

$$(6.2.1) \quad \langle h_{e_1} \otimes \dots \otimes h_{e_{n-1}} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_r]}$$

where $e_1, \dots, e_{n-1} \in \mu_r$ are group elements of age 1, and e_n is arbitrary, and also the only element for which we allow insertion of a ψ -class. (This implies that $\delta_T^{(1)} + \delta_T^{(2)} + \delta_T^{(3)} = |T|$ for all $T \subset [n-1]$.)

We want to determine the integral of a summand on the right-hand-side of equation (6.1.2) related to a partition \mathcal{P} of $[n-1]$, after inserting an additional ψ -class at the n -th marked point. Let T_1, \dots, T_k be the elements of $\mathcal{P}_{\geq 2}$.

$$\psi_n^\nu \cdot c(\mathcal{P}) = \psi_n^\nu \cdot \prod_{i=1}^k x_{T_i} \prod_{\substack{T \subseteq [n-1]/(\mathcal{P}) \\ 2 \leq |T|}} (1 + x_T) \prod_{a=1}^N \prod_{p=\langle \delta_{[n-1]}^{(a)} \rangle}^{\delta_{[n-1]}^{(a)}-1} (t_a - p\psi_n)$$

This term is supported on the intersection $D^{T_1} \cap \dots \cap D^{T_k}$, isomorphic to $\overline{M}_{0, T_1 \cup \{*\}} \times \dots \times \overline{M}_{0, T_k \cup \{*\}} \times \overline{M}_{0, [n]/(\mathcal{P})}$; so in order to determine the integral of $c(\mathcal{P}) \cdot \psi_n^\nu$, we will write it as a product of $D^{T_1} \dots D^{T_k}$ with factors that are pulled back from one of the components above.

The numerator of x_{T_j} is the only factor that has terms coming from $\overline{M}_{0, T_j \cup \{*\}}$, while its denominator involves ψ_{T_j} , which is the ψ -class of the node corresponding to the marking T_j on $\overline{M}_{0, [n]/(\mathcal{P})}$. To examine the numerator more closely, we first factor out D^{T_j} :

$$\begin{aligned} \prod_{a,p} (1 - t_a^{-1} p\psi_{T_j^C}) - \prod_{a,p} (1 + t_a^{-1} p\psi_{T_j}) &= \sum_{k \geq 0} \beta_k ((-1)^k \psi_{T_j^C}^k - \psi_{T_j}^k) \\ &= D^{T_j} \cdot \sum_{k \geq 0} \beta_k \sum_{\ell=0}^{k-1} (-\psi_{T_j^C})^\ell \psi_{T_j}^{k-1-\ell}. \end{aligned}$$

The largest power of $\psi_{T_j^C}$ which appears in the last expression is

$$\sum_{a=1}^3 (\delta_{T_j}^{(a)} - \langle \delta_{T_j}^{(a)} \rangle) - 1 = |T_j| - 1 - \sum_{a=1}^3 \langle \delta_{T_j}^{(a)} \rangle.$$

As the dimension of $\overline{M}_{0, T_j \cup \{*\}}$ is $|T_j| - 2$, the expression only has a term in the top degree if $\sum_{a=1}^3 \langle \delta_{T_j}^{(a)} \rangle = 1$, which means that $e_{T_j} = \prod_{i \in T_j} e_i$ acts with age 1 on \mathbb{C}^3 and acts nontrivially in each coordinate direction. By the balancing condition, e_{T_j} is prescribing the monodromy of the node as seen from the component corresponding to $[n]/(\mathcal{P})$. As the integral of $\psi_{T_j^C}^{|T_j|-2}$ is one, the integral of the above product

on $\overline{M}_{0, T_j \cup \{*\}}$ is $\prod_{a=1}^3 t_a^{-\lceil \delta_{T_j}^{(a)} \rceil} (\delta_{T_j}^{(a)} - 1)!$ if the condition on e_{T_j} is satisfied, and 0 otherwise.

On $\overline{M}_{0, [n]/(\mathcal{P})}$, we are left with the following product:

$$\psi_n^\nu \cdot \prod_{a=1}^3 t_a^{-1 + \delta_{[n]}^{(a)}} \prod_{\emptyset \neq T \subseteq [n-1]/(\mathcal{P})} \prod_{p=\langle \delta_T^{(a)} \rangle}^{\delta_T^{(a)}-1} \frac{t_a + p\psi_T + pD^T}{t_a + p\psi_T}$$

Here we used the same conventions as for formula (5.2.1), applied to the set $[n-1]/(\mathcal{P})$ (so for example $D^{\{T_j\}}$ is identified with $-\psi_{\{T_j\}}$, which is the ψ -class

of a single marking; the term related to T_j in the above product is the denominator of x_{T_j}); and we extended $\delta_T^{(a)}$ in the obvious way from subsets of $[n-1]$ to subsets of the quotient set $[n-1]/(\mathcal{P})$. Now by the remarks in section 5.2, based on lemma A.1.1, this product computes the Chern class of the obstruction bundle on $\overline{M}_{0,[n]/(\mathcal{P})}((e_i)_{i \in [n]/(\mathcal{P})}; B\mu_r)$. Its integral is thus given by the equivariant Gromov-Witten invariant

$$\left\langle \bigotimes_{i \in [n-1]/(\mathcal{P})} h_{e_i} \otimes \psi_n^\nu h_{e_n} \right\rangle_{0,[n]/(\mathcal{P})}^{[\mathbb{C}^3/\mu_r]}.$$

This proves the following recursion:

Proposition 6.2.1. *For an equivariant Gromov-Witten invariant of $[\mathbb{C}^3/\mu_r]$ as in equation (6.2.1) with the assumptions above, let S be the set of non-trivial partitions \mathcal{P} of $[n-1]$ such that for every $T \in \mathcal{P}_{\geq 2}$, the group element $e_{T_j} = \prod_{i \in T_j} e_i$ acts with age 1 on \mathbb{C}^3 , and non-trivial in every coordinate direction. Then the following recursive formula holds:*

$$\begin{aligned} \langle h_{e_1} \otimes \cdots \otimes h_{e_{n-1}} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_r]} &= \langle h_{e_1} \otimes \cdots \otimes h_{e_{n-1}} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_r]; \text{weighted}} + \\ &+ \sum_{\mathcal{P} \in S} (-1)^{|\mathcal{P}_{\geq 2}|+1} \prod_{T \in \mathcal{P}_{\geq 2}} \prod_{a=1}^3 (\delta_T^{(a)} - 1)! \left\langle \bigotimes_{i \in [n-1]/(\mathcal{P})} h_{e_i} \otimes \psi_n^\nu h_{e_n} \right\rangle_{0,[n]/(\mathcal{P})}^{[\mathbb{C}^3/\mu_r]}. \end{aligned}$$

Here the invariant with superscript “weighted” means the invariant as computed by using the moduli space of weighted stable maps instead of the ordinary moduli space, with weight data chosen as in section 3.4. These invariants are given, up to a multiplication with a monomial in the t_a , by the $(n-3-\nu)$ -th elementary symmetric function of the variables $t_a^{-1}(\delta_{[n-1]}^{(a)} - 1), t_a^{-1}(\delta_{[n-1]}^{(a)} - 2), \dots, t_a^{-1}(\delta_{[n-1]}^{(a)})$ for $a = 1, 2, 3$.

6.3. Recursions for $[\mathbb{C}^3/\mu_3]$. The recursion of proposition 6.2.1 simplifies further in the case $[\mathbb{C}^3/\mu_3]$ for the diagonal representation of μ_3 . The only group element of age 1 is $\omega = e^{\frac{2\pi i}{3}}$. We have $\delta_T^{(a)} = \frac{|T|}{3}$ for all $T \subset [n-1]$ and $a = 1, 2, 3$. The set S contains the partitions \mathcal{P} of $[n-1]$ so that every $T \in \mathcal{P}$ has size $3m_T + 1$ for some $m_T \in \mathbb{Z}_{\geq 0}$. The summand for \mathcal{P} in the formula of proposition 6.2.1 depends only on the sizes of the subsets, not on the actual subsets; if we set $p = \sum_T m_T$, we can thus reduce the above sum to a sum over partitions $\underline{m} = (m_1, \dots, m_k)$ of p , for all $p \geq 1$ with $n-3p \geq 3$. For any such partition, let $M(n-1, \underline{m})$ be the multinomial coefficient

$$M(n-1, \underline{m}) = \binom{n-1}{3m_1+1, \dots, 3m_k+1, n-1-\sum_j (3m_j+1)}$$

counting the ways to distribute $n-1$ markings on the different components.

Proposition 6.3.1.

$$\langle h_\omega^{\otimes n-1} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_3]} = \langle h_\omega^{\otimes n-1} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_3]; \text{weighted}}$$

$$+ \sum_{p=1}^{\lfloor \frac{n-3}{3} \rfloor} \sum_{\underline{m}} \frac{(-1)^{|\underline{m}|+1}}{|\text{Aut } \underline{m}|} \prod_{j=1}^k \left((m_j - \frac{2}{3})! \right)^3 M(n-1, \underline{m}) \langle h_\omega^{\otimes n-1-3p} \otimes \psi_n^\nu h_{e_n} \rangle_{0,n}^{[\mathbb{C}^3/\mu_3]}$$

The superscript “weighted” means the same as before; if $e_n = \omega$ and $\nu = 0$, this weighted Gromov-Witten invariant is just given as $(-1)^{n+1} \left((\frac{n-4}{3})! \right)^3 \frac{1}{3}$; otherwise it is an elementary symmetric function.




FIGURE 1. Comb for $n = 30$, $p = 7$ and $\underline{m} = (1, 1, 2, 3)$

A maple program implementing some of these recursions is available from the authors upon request. The numbers match the calculations of [ABK08], [CCIT07a] and [CC07].

As the recursions are linear, it is not hard to invert the matrix and obtain a direct formula. Let $I_\ell = \langle h_\omega^{\otimes 3\ell+3} \rangle_{0,3\ell+3}^{[\mathbb{C}^3/\mu_3]}$. Proposition 6.3.1 implies that

$$\sum_{p=0}^{\ell} (-1)^p C_{p,\ell} I_{\ell-p} = (-1)^\ell \left(\left(\ell - \frac{1}{3} \right)! \right)^3 \frac{1}{3},$$

where $C_{0,\ell} = 1$ and for $0 < p \leq \ell$, $C_{p,\ell}$ is the sum over partitions $\underline{m} = (m_1, \dots, m_k)$ of p , with $k \leq 3(\ell - p) + 2$, of the quantity

$$\frac{1}{|\text{Aut } \underline{m}|} \prod_{j=1}^k \left((m_j - \frac{2}{3})! \right)^3 \cdot \binom{3\ell + 2}{3m_1 + 1, \dots, 3m_k + 1, 3(\ell - p) + 2 - k}.$$

Let $D_{p,\ell} = C_{\ell-p,\ell}$. By inverting the matrix we obtain the formula:

$$3(-1)^\ell I_\ell = \sum_{\substack{S \subseteq [0, \ell-1] \\ S = \{x_0, \dots, x_q\}}} (-1)^{|S|} \left(\left(x_0 - \frac{1}{3} \right)! \right)^3 D_{x_0, x_1} \cdots D_{x_{q-1}, x_q} D_{x_q, \ell},$$

where $[0, \ell - 1] = \{0, 1, \dots, \ell - 1\}$ and it is assumed that $x_0 < x_1 < \dots < x_q$. For $S = \emptyset$, the summand is taken to be $((\ell - 1/3)!)^3$.

6.4. Inversion of the “mirror map”. The recursion in this case can also be derived from the results in [CCIT07a]. We explained above in section 3.5 that their I -function $I^{\text{tw}}(t)$ is almost identical (up to high powers of z) to the “weighted J -function” $J^{X;\text{weighted}}$.

From general principles of Givental’s formalism they deduce that for the coordinate change $\tau(t): H \rightarrow H$, called the “mirror map”, given by

$$\tau(t_0 h_1 + t_1 h_\omega) = t_0 h_1 + \sum_{k \geq 0} \frac{(-1)^{3k} (t_1)^{3k+1}}{(3k+1)!} \left((k - \frac{2}{3})! \right)^3 h_\omega$$

the twisted I -function and the twisted J -function can be related after setting the dual coordinate of h_{ω^2} equal to zero:

$$I^{\text{tw}}(t_0 h_1 + t_1 h_\omega, z) = J^X(\tau(t_0 h_1 + t_1 h_\omega), z)$$

Our recursive formula can be recovered by comparing coefficients of these two power series; in other words, the sum over partitions \mathcal{P} in proposition 6.3.1 is a combinatorial inversion of the mirror map $\tau(t)$.

More precisely, given any two power series $A(t_0 h_1 + t_1 h_\omega, z)$, $B(t_0 h_1 + t_1 h_\omega, z)$ related by

$$A(t_0 h_1 + t_1 h_\omega, z) = B(\tau(t_0 h_1 + t_1 h_\omega), z)$$

we can recursively recover the coefficients of B by comparing coefficients of powers of t_1 . If we write $A(t_1 h_1) = \sum_k \frac{a_k}{k!} t_1^k$ and $B(t_1 h_1) = \sum_k \frac{b_k}{k!} t_1^k$ with $a_k, b_k \in H[[z^{-1}]]$, the recursion will look exactly as proposition 6.3.1 with the “weighted invariants” replaced by a_k and the actual invariants replaced by b_k . In particular, setting A to the identity power series $A(t_0 h_1 + t_1 h_\omega) = t_0 h_1 + t_1 h_\omega$ yields an inversion of the mirror map that can also be interpreted as a sum over comb curves.

6.5. Equivariant recursions. The methods of this section are sufficient to produce a linear recursion for the equivariant descendant Gromov-Witten invariants of $[\mathbb{C}^N / \mu_r]$. However, this requires one to allow ψ -classes at every marked point. As in section 6.2, one can use lemma 6.1.1 to expand the equivariant Euler class of the obstruction bundle, and to each partition of $[n-1]$ one should associate a comb as before, where the n -th marked point is on the head of the comb. For each tooth of the comb, we can write the numerator of x_T as

$$D^T \cdot \sum_{k > 0} \beta_k \sum_{\ell=0}^{k-1} (-\psi_{TC})^\ell \psi_T^{k-1-\ell},$$

just as in section 6.2. The exponent of ψ_{TC} which leads to a nonzero integral is determined by the descendant exponents chosen for the marked points in T . The integral over the tooth can then be computed using the well-known formula

$$\int_{\overline{M}_{0,n}} \psi_1^{a_1} \cdots \psi_n^{a_n} = \binom{n-3}{a_1, \dots, a_n}.$$

We are left with a polynomial in ψ_T , the ψ -class of the node on the main component (and hence our method does not yield a recursion for non-descendant invariants only).

In summary, for each partition of $[n-1]$, one gets a linear combination of equivariant Gromov-Witten invariants with fewer marked points, each with a combinatorial factor. These must be summed together and added to the weighted invariant, just as in section 6.2.

APPENDIX A. COMBINATORICS OF DIVISORS ON $\overline{M}_{0,n}$

A.1. Notations for divisors. This section reviews notations for divisors on $\overline{M}_{0,n}$, some of which is introduced in this paper. The standard relations are reviewed, and a combinatorial proof of a key simplifying relation, used in section 6, is worked out.

First recall the vital divisor D^T , introduced by Keel [Kee92], where T is any subset of $[n]$ having at least 2 and at most $n-2$ elements. This divisor is the locus of curves having a node which separates the markings into T and T^C . Here complements are always taken within $[n]$. To make this into a correspondence, assume throughout this appendix that S and T are subsets of $[n-1]$. It is natural to define $D^{[n-1]}$ to be $-\psi_n$, which comes from the work of de Concini and Procesi [DCP95]. From their point of view, $D^{[n-1]}$ is the pullback of minus the hyperplane class under a sequence of blowups producing $\overline{M}_{0,n}$ from \mathbb{P}^{n-3} . This sequence of blowups is the same one discussed in section 3.2 in the context of weighted stable maps. Under this blowup description of $\overline{M}_{0,n}$, D^T is the exceptional divisor of the blowup in the proper transform of the linear space generated by the points labeled by $[n-1] \setminus T$.

The ring $H^*(\overline{M}_{0,n})$ is generated by the divisors D^T for $T \subseteq [n-1]$, $|T| \geq 2$, with relations given by

$$(A.1.1) \quad D^S D^T \text{ if } S \text{ and } T \text{ are incomparable and } S \cap T \neq \emptyset,$$

$$(A.1.2) \quad \sum_{i,j \in T} D^T \text{ for every } i \neq j \in [n-1] \text{ [DCP95].}$$

Geometrically, the first relation is due to the fact that the exceptional divisors D^S and D^T are disjoint, and the second is due to the fact that the preimage of the hyperplane in \mathbb{P}^{n-3} generated by all points except i and j consists of the proper transform $D^{i,j}$ together with all the exceptional divisors D^T for $i, j \in T$. Fix $T \subsetneq [n-1]$ and choose $i \notin T$ and $j \in T$. Then relations (A.1.1, A.1.2) imply that

$$(A.1.3) \quad D^T \sum_{\substack{S: i \in S \\ S \supset T}} D^S = 0.$$

For any nonempty subset $T \subseteq [n-1]$, we introduce the notation

$$(A.1.4) \quad \psi_T := \sum_{S \supsetneq T} D^S.$$

If $T = \{i\}$, then this recovers the ψ -class at the i -th marked point. Indeed, we have for any distinct j, k different from i (recalling the convention $n \notin T$),

$$\psi_i = \sum_{\substack{j,k \in T \\ i \notin T}} D^T + \sum_{\substack{i \in T \\ j,k \notin T}} D^T = - \sum_{i,j,k \in T} D^T + \sum_{\substack{i \in T \\ j,k \notin T}} D^T + \sum_{i,j \in T} D^T + \sum_{i,k \in T} D^T = \psi_{\{i\}},$$

the last equality following from an inclusion-exclusion argument. By definition, $\psi_{[n-1]} = 0$, and we also set $D^i = -\psi_i$ for $1 \leq i \leq n-1$. Finally, we define

$$\psi_{T^C} := -D^T - \psi_T$$

for $T \subseteq [n-1]$, which we found useful in section 6.

The Chern class formula of theorem 5.1.1 can now be expressed as

$$(A.1.5) \quad c = \prod_{\emptyset \neq T \subseteq [n-1]} \prod_{p=\langle \delta_T \rangle}^{\delta_T-1} \left(\frac{1 + p(D^T + \psi_T)}{1 + p\psi_T} \right),$$

where $\delta_T = \sum_{i \in T} \delta_i$ and δ_i are chosen so that $0 \leq \delta_i < 1$. The following lemma shows that if every $\delta_i \geq 0$, this expression is periodic in each δ_i with period 1; so it defines a continuous, piecewise-analytic function from an $(n-1)$ -dimensional real torus into the cohomology of $\overline{M}_{0,n}$ with real coefficients. This is used in section 6 to produce recursions for the Gromov-Witten invariants.

Lemma A.1.1. *Let $\delta_1, \dots, \delta_{n-1}$ be real numbers, and for any subset $T \subseteq [n-1]$, define $\delta_T = \sum_{i \in T} \delta_i$. Then*

$$(A.1.6) \quad \prod_{1 \in T} \frac{1 + \delta_T(D^T + \psi_T)}{1 + \delta_T\psi_T} = 1.$$

Proof. For $1 \leq k \leq n-1$, let

$$E_k = \sum_{i=2}^{n-1} \delta_i \sum_{\substack{1, i \in T \\ |T| > k}} D^T + \delta_1 \sum_{\substack{1 \in T \\ |T| > k}} D^T,$$

and let

$$A_k = (1 + E_k) \prod_{\substack{1 \in T \subseteq [n-1] \\ |T| \leq k}} \frac{1 + \delta_T(D^T + \psi_T)}{1 + \delta_T\psi_T}.$$

Then it must be shown that $A_{n-1} = 1$. As $E_{n-2} = \delta_{[n-1]} D^{[n-1]}$, it follows that $A_{n-2} = A_{n-1}$. Moreover, $E_1 = \delta_1 \psi_1$, so $A_1 = 1$. It remains to show that $A_k = A_{k-1}$ for $1 < k < n-1$.

Note that for any divisors x, y, z ,

$$(A.1.7) \quad \frac{(1+x)(1+y)}{1+z} = 1 + x + y - z \quad \text{if } (x-z)(y-z) = 0.$$

Fix T with $1 \in T$ and $|T| = k$, and let x be any expression of the form $E_k + \sum_{S \in \sigma} \alpha_S D^S$, where σ is a collection of subsets $S \subseteq [n-1]$ with $|S| = k$ and

$1 \in S \neq T$. Then $D^T(x - \delta_T \psi_T) = 0$ by the following argument. For any $S \in \sigma$, $1 \in S \cap T$ and $|S| = |T|$, so $D^S D^T = 0$. Moreover,

$$E_k - \delta_T \psi_T = \sum_{i: i \notin T} \delta_i \sum_{\substack{S: i \in S \\ T \subseteq S}} D^S + \sum_{\substack{|S| > k \\ S \not\supseteq T \\ 1 \in S}} \delta_S D^S.$$

Since D^T annihilates the first term on the right hand side by (A.1.3) and annihilates the second by (A.1.1), it follows that $D^T(E_k - \delta_T \psi_T) = 0$, completing the verification of the claim.

It follows that one can iterate through all sets T with $|T| = k$ to eliminate those factors from the expression for A_k and apply relation (A.1.7) at each step. Since

$$E_k + \sum_{\substack{1 \in T \\ |T|=k}} \delta_T D^T = E_{k-1},$$

it follows that $A_k = A_{k-1}$, completing the proof. \square

If in (A.1.5), one were to add 1 to δ_1 (and thus add one to each δ_T for $1 \in T$) then one would be multiplying c by the left hand side of (A.1.6). Therefore c does not change after the translation $\delta_i \mapsto \delta_i + 1$. Using the notational conventions of section 1.5, the formula for c makes sense for negative values of δ_i , and the same argument shows that it remains invariant under integer translations. It seems natural to regard $\delta_1, \dots, \delta_n$ as coordinates on $(\mathbb{R}/\mathbb{Z})^n$ satisfying $\sum_{i=1}^n \delta_i = 0$.

A.2. Restricting to D^T . It is a standard fact for any proper subset $T \subset [n-1]$ containing at least two elements, $D^T \cong \overline{M}_{0,|T|+1} \times \overline{M}_{0,n-|T|+1}$, with the node counting as an extra marked point on each factor. The restrictions of divisors D^S and ψ_S to D^T are easily computed if one uses subsets of T for the divisors on the first factor on subsets of the quotient set $[n-1]/T$ for divisors on the second factor. So on the first factor, the node counts as the extra marked point that all sets must avoid. For sets $S \subset T$, we use D_*^S and ψ_S^* to notate divisors on $\overline{M}_{0,|T|+1}$ and we likewise use D_\bullet^S and ψ_S^\bullet for divisors on $\overline{M}_{0,n-|T|+1}$. Now the following formulas hold for any nonempty set $S \subseteq [n-1]$.

$$(A.2.1) \quad D^S|_{D^T} = \begin{cases} D_*^S \otimes 1, & \text{if } S \subsetneq T \\ 1 \otimes D_\bullet^S, & \text{if } S \cap T = \emptyset \\ 1 \otimes D_\bullet^{S/T}, & \text{if } S \supsetneq T \\ D_*^T \otimes 1 + 1 \otimes D_\bullet^{T/T}, & \text{if } S = T \\ 0, & \text{otherwise} \end{cases}$$

$$(A.2.2) \quad \psi_S|_{D^T} = \begin{cases} \psi_S^* \otimes 1, & \text{if } S \subsetneq T \\ 1 \otimes \psi_S^\bullet, & \text{if } S \cap T = \emptyset \\ 1 \otimes \psi_{S/T}^\bullet, & \text{if } S \supseteq T \end{cases}$$

Note that the fourth line of (A.2.1) is another way of writing the standard fact that the restriction of $-D^T$ to D^T is the sum of the ψ -classes at the node on the two

components. Moreover, the third line of (A.2.2) shows that ψ_T restricts to D^T as the ψ class of the node on the component corresponding to T^C . So our definition of ψ_T^C as $-D^T - \psi_T$ ensures that ψ_T^C restricts to the ψ class of the node on the other component.

REFERENCES

- [ABK08] Mina Aganagic, Vincent Bouchard, and Albrecht Klemm. Topological strings and (almost) modular forms. *Comm. Math. Phys.*, 277(3):771–819, 2008, arXiv:hep-th/0607100.
- [Abr05] Dan Abramovich. Lectures on Gromov-Witten invariants of orbifolds, 2005, arXiv:math.AG/0512372.
- [ACV03] Dan Abramovich, Alessio Corti, and Angelo Vistoli. Twisted bundles and admissible covers. *Comm. Algebra*, 31(8):3547–3618, 2003, arXiv:math.AG/0106211. Special issue in honor of Steven L. Kleiman.
- [AG08] Valery Alexeev and G. Michael Guy. Moduli of weighted stable maps and their gravitational descendants. *J. Inst. Math. Jussieu*, 7(3):425–456, 2008, arXiv:math.AG/0607683.
- [AV02] Dan Abramovich and Angelo Vistoli. Compactifying the space of stable maps. *J. Amer. Math. Soc.*, 15(1):27–75 (electronic), 2002, arXiv:math.AG/9908167.
- [BG06] Jim Bryan and Tom Graber. The Crepant Resolution Conjecture, 2006, arXiv:math.AG/0610129.
- [BGP08] Jim Bryan, Tom Graber, and Rahul Pandharipande. The orbifold quantum cohomology of $\mathbb{C}^2/\mathbb{Z}_3$ and Hurwitz-Hodge integrals. *J. Algebraic Geom.*, 17(1):1–28, 2008, arXiv:math.AG/0510335.
- [BM06] Arend Bayer and Yuri I. Manin. Stability Conditions, Wall-crossing and weighted Gromov-Witten Invariants, 2006, arXiv:math.AG/0607580. To appear in Moscow Math Journal.
- [Cad07] Charles Cadman. Using stacks to impose tangency conditions on curves. *Amer. J. Math.*, 129(2):405–427, 2007.
- [CC07] Charles Cadman and Renzo Cavalieri. Gerby localization, Hodge integrals, and the GW theory of $[\mathbb{C}^3/\mathbb{Z}_3]$, 2007, arXiv:0705.2158.
- [CC08] Charles Cadman and Linda Chen. Enumeration of rational plane curves tangent to a smooth cubic. *Adv. Math.*, 219(1):316–343, 2008, arXiv:math.AG/0701406.
- [CCIT06] Tom Coates, Alessio Corti, Hiroshi Iritani, and Hsian-Hua Tseng. Wall-Crossings in Toric Gromov-Witten Theory I: Crepant Examples, 2006, arXiv:math.AG/0611550.
- [CCIT07a] Tom Coates, Alessio Corti, Hiroshi Iritani, and Hsian-Hua Tseng. Computing Genus-Zero Twisted Gromov-Witten Invariants, 2007, arXiv:math.AG/0702234.
- [CCIT07b] Tom Coates, Alessio Corti, Hiroshi Iritani, and Hsian-Hua Tseng. The crepant resolution conjecture for type A surface singularities, 2007, arXiv:0704.2034.
- [CG07] Tom Coates and Alexander Givental. Quantum Riemann-Roch, Lefschetz and Serre. *Ann. of Math. (2)*, 165(1):15–53, 2007, arXiv:math.AG/0110142.
- [DCP95] C. De Concini and C. Procesi. Hyperplane arrangements and holonomy equations. *Selecta Math. (N.S.)*, 1(3):495–535, 1995.
- [FP00] C. Faber and R. Pandharipande. Logarithmic series and Hodge integrals in the tautological ring. *Michigan Math. J.*, 48:215–252, 2000, arXiv:math.AG/0002112. With an appendix by Don Zagier, Dedicated to William Fulton on the occasion of his 60th birthday.
- [Giv01] Alexander B. Givental. Gromov-Witten invariants and quantization of quadratic Hamiltonians. *Mosc. Math. J.*, 1(4):551–568, 645, 2001, arXiv:math.AG/0108100. Dedicated to the memory of I. G. Petrovskii on the occasion of his 100th anniversary.
- [Giv04] Alexander B. Givental. Symplectic geometry of Frobenius structures. In *Frobenius manifolds*, Aspects Math., E36, pages 91–112. Vieweg, Wiesbaden, 2004, arXiv:math.AG/0305409.

- [Has03] Brendan Hassett. Moduli spaces of weighted pointed stable curves. *Adv. Math.*, 173(2):316–352, 2003, arXiv:math.AG/0205009.
- [JK02] Tyler J. Jarvis and Takashi Kimura. Orbifold quantum cohomology of the classifying space of a finite group. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 123–134. Amer. Math. Soc., Providence, RI, 2002, arXiv:math.AG/0112037.
- [Kee92] Sean Keel. Intersection theory of moduli space of stable n -pointed curves of genus zero. *Trans. Amer. Math. Soc.*, 330(2):545–574, 1992.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.
- [MM08] Anca M. Mustață and Andrei Mustață. The Chow ring of $\overline{M}_{0,m}(\mathbb{P}^n, d)$. *J. Reine Angew. Math.*, 615:93–119, 2008, arXiv:math.AG/0507464.
- [Rua02] Yongbin Ruan. Stringy geometry and topology of orbifolds. In *Symposium in Honor of C. H. Clemens (Salt Lake City, UT, 2000)*, volume 312 of *Contemp. Math.*, pages 187–233. Amer. Math. Soc., Providence, RI, 2002.
- [Rua06] Yongbin Ruan. The cohomology ring of crepant resolutions of orbifolds. In *Gromov-Witten theory of spin curves and orbifolds*, volume 403 of *Contemp. Math.*, pages 117–126. Amer. Math. Soc., Providence, RI, 2006, arXiv:math.AG/0108195.
- [SGA71] *Théorie des intersections et théorème de Riemann-Roch*. Springer-Verlag, Berlin, 1971. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6), Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Avec la collaboration de D. Ferrand, J. P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J. P. Serre, Lecture Notes in Mathematics, Vol. 225.
- [Tse05] Hsian-Hua Tseng. Orbifold Quantum Riemann-Roch, Lefschetz and Serre, 2005, arXiv:math.AG/0506111.

UNIVERSITY OF UTAH, DEPARTMENT OF MATHEMATICS, 155 SOUTH 1400 EAST, ROOM 233, SALT LAKE CITY, UT 84112

E-mail address: bayer@math.utah.edu

UNIVERSITY OF BRITISH COLUMBIA, DEPARTMENT OF MATHEMATICS, 1984 MATHEMATICS RD, VANCOUVER, BC, CANADA V6P 4M8

E-mail address: cadman@math.ubc.ca